

Protocol

Automated Percutaneous and Percutaneous Endoscopic Discectomy

(70118)

Medical Benefit		Effective Date: 10/01/18	Next Review Date: 07/19
Preauthorization	No	Review Dates: 02/07, 01/08, 01/09, 01/10, 01/11, 01/12, 01/13, 09/13, 07/14, 07/15, 07/16, 07/17, 07/18	

This protocol considers this test or procedure investigational. If the physician feels this service is medically necessary, preauthorization is recommended.

The following protocol contains medical necessity criteria that apply for this service. The criteria are also applicable to services provided in the local Medicare Advantage operating area for those members, unless separate Medicare Advantage criteria are indicated. If the criteria are not met, reimbursement will be denied and the patient cannot be billed. Please note that payment for covered services is subject to eligibility and the limitations noted in the patient's contract at the time the services are rendered.

Populations	Interventions	Comparators	Outcomes
Individuals: <ul style="list-style-type: none">• With herniated intervertebral disc(s)	Interventions of interest are: <ul style="list-style-type: none">• Automated percutaneous discectomy	Comparators of interest are: <ul style="list-style-type: none">• Conservative therapy• Open discectomy or microdiscectomy	Relevant outcomes include: <ul style="list-style-type: none">• Symptoms• Functional outcomes• Quality of life• Treatment-related morbidity
Individuals: <ul style="list-style-type: none">• With herniated intervertebral disc(s)	Interventions of interest are: <ul style="list-style-type: none">• Percutaneous endoscopic discectomy	Comparators of interest are: <ul style="list-style-type: none">• Conservative therapy• Open discectomy or microdiscectomy	Relevant outcomes include: <ul style="list-style-type: none">• Symptoms• Functional outcomes• Quality of life• Treatment-related morbidity

DESCRIPTION

Surgical management of herniated intervertebral discs most commonly involves discectomy or microdiscectomy, performed manually through an open incision. Automated percutaneous discectomy involves placement of a probe within the intervertebral disc under image guidance with aspiration of disc material using a suction cutting device. Removal of disc herniations under endoscopic visualization is also being investigated. Endoscopic discectomy involves the percutaneous placement of a working channel under image guidance, followed by visualization of the working space and instruments through an endoscope, and aspiration of disc material.

SUMMARY OF EVIDENCE

For individuals who have herniated intervertebral disc(s) who receive automated percutaneous discectomy, the evidence includes randomized controlled trials (RCTs) and systematic reviews of RCTs. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. The published evidence from small RCTs is insufficient to evaluate the impact of automated percutaneous discectomy on net health outcomes. Well-designed and executed RCTs are needed to determine the benefits and risks of this procedure. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have herniated intervertebral disc(s) who receive percutaneous endoscopic discectomy, the evidence includes a number of RCTs and systematic reviews of RCTs. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. Many of the RCTs were conducted at a single center in Europe. Some trials have reported outcomes at least as good as traditional approaches with an open incision, while one RCT from a different center in Europe reported a trend toward increased complications and reherniations using an endoscopic approach. There are few reports from the United States. Results from a number of moderately large ongoing RCTs are anticipated in the next two to three years. The evidence is insufficient to determine the effects of the technology on health outcomes.

POLICY

Automated percutaneous discectomy is considered **investigational** as a technique of intervertebral disc decompression in patients with back pain and/or radiculopathy related to disc herniation in the lumbar, thoracic, or cervical spine.

Percutaneous endoscopic discectomy is considered **investigational** as a technique of intervertebral disc decompression in patients with back pain and/or radiculopathy related to disc herniation in the lumbar, thoracic, or cervical spine.

MEDICARE ADVANTAGE

For Medicare Advantage percutaneous image guided lumbar decompression (PILD) may have potential for coverage when provided through Coverage with Evidence Development (CED) for members with lumbar spinal stenosis who meet the criteria of and are enrolled in an approved clinical study.

Effective for services performed on or after December 7, 2016, there may be a potential for coverage through a prospective, longitudinal study of PILD procedures using an FDA-approved/cleared device that completed a CMS-approved randomized control trial (RCT) that met CMS criteria.

MEDICARE ADVANTAGE POLICY GUIDELINES

Approved studies will be identified on the CMS website - <https://www.cms.gov/Medicare/Coverage/Coverage-with-Evidence-Development/index.html>.

BACKGROUND

Back pain or radiculopathy related to herniated discs is an extremely common condition and a frequent cause of chronic disability. Although many cases of acute low back pain and radiculopathy will resolve with conservative care, surgical decompression is often considered when the pain is unimproved after several months and is clearly neuropathic in origin, resulting from irritation of the nerve roots. Open surgical treatment typically consists of discectomy in which the extruding disc material is excised. When performed with an operating microscope, the procedure is known as microdiscectomy.

Minimally invasive options have also been researched, in which some portion of the disc is removed or ablated, although these techniques are not precisely targeted at the offending extruding disc material. Ablative techniques include laser discectomy and radiofrequency decompression (see the Decompression of the Intervertebral Disc Using Laser Energy [Laser Discectomy] or Radiofrequency Coblation [Nucleoplasty] Protocol). Intradiscal electrothermal annuloplasty is another minimally invasive approach to low back pain. In this technique,

radiofrequency energy is used to treat the surrounding disc annulus (see the Percutaneous Intradiscal Electrothermal Annuloplasty, Radiofrequency Annuloplasty, and Biacuplasty Protocol).

Herein we address automated percutaneous and endoscopic discectomy, in which the disc decompression is accomplished by the physical removal of disc material rather than its ablation. Traditionally, discectomy was performed manually through an open incision, using cutting forceps to remove nuclear material from within the disc annulus. This technique was modified by automated devices that involve placement of a probe within the intervertebral disc and aspiration of disc material using a suction cutting device. Endoscopic techniques may be intradiscal or may involve extraction of noncontained and sequestered disc fragments from inside the spinal canal using an interlaminar or transforaminal approach. Following insertion of the endoscope, decompression is performed under visual control.

REGULATORY STATUS

The Dekompressor® Percutaneous Discectomy Probe (Stryker), Herniatome Percutaneous Discectomy Device (Gallini Medical Devices), and the Nucleotome® (Clarus Medical) are examples of percutaneous discectomy devices that have been cleared for marketing by the U.S. Food and Drug Administration (FDA) through the 510(k) process. The FDA indication for these products is for “aspiration of disc material during percutaneous discectomies in the lumbar, thoracic and cervical regions of the spine.” FDA product code: HRX.

A variety of endoscopes and associated surgical instruments have also been cleared for marketing by FDA through the 510(k) process.

RELATED PROTOCOLS

Decompression of the Intervertebral Disc Using Laser Energy (Laser Discectomy) or Radiofrequency Coblation (Nucleoplasty)

Percutaneous Intradiscal Electrothermal Annuloplasty, Radiofrequency Annuloplasty, and Biacuplasty

Services that are the subject of a clinical trial do not meet our Technology Assessment Protocol criteria and are considered investigational. *For explanation of experimental and investigational, please refer to the Technology Assessment Protocol.*

It is expected that only appropriate and medically necessary services will be rendered. We reserve the right to conduct prepayment and postpayment reviews to assess the medical appropriateness of the above-referenced procedures. **Some of this protocol may not pertain to the patients you provide care to, as it may relate to products that are not available in your geographic area.**

REFERENCES

We are not responsible for the continuing viability of web site addresses that may be listed in any references below.

1. Rasouli MR, Rahimi-Movaghar V, Shokrane F, et al. Minimally invasive discectomy versus microdiscectomy/ open discectomy for symptomatic lumbar disc herniation. *Cochrane Database Syst Rev.* 2014;9:CD010328. PMID 25184502

2. Lewis RA, Williams NH, Sutton AJ, et al. Comparative clinical effectiveness of management strategies for sciatica: systematic review and network meta-analyses. *Spine J.* Jun 1 2015;15(6):1461-1477. PMID 24412033
3. Manchikanti L, Singh V, Falco FJ, et al. An updated review of automated percutaneous mechanical lumbar discectomy for the contained herniated lumbar disc. *Pain Physician.* Apr 2013;16(2 Suppl):SE151-184. PMID 23615890
4. Manchikanti L, Singh V, Calodney AK, et al. Percutaneous lumbar mechanical disc decompression utilizing Dekompressor(R): an update of current evidence. *Pain Physician.* Apr 2013;16(2 Suppl):SE1-24. PMID 23615884
5. Vorobeychik Y, Gordin V, Fuzaylov D, et al. Percutaneous mechanical disc decompression using Dekompressor device: an appraisal of the current literature. *Pain Med.* May 2012;13(5):640-646. PMID 22494347
6. Singh V, Benyamin RM, Datta S, et al. Systematic review of percutaneous lumbar mechanical disc decompression utilizing Dekompressor. *Pain Physician.* May-Jun 2009;12(3):589-599. PMID 19461825
7. Hirsch JA, Singh V, Falco FJ, et al. Automated percutaneous lumbar discectomy for the contained herniated lumbar disc: a systematic assessment of evidence. *Pain Physician.* May-Jun 2009;12(3):601-620. PMID 19461826
8. Revel M, Payan C, Vallee C, et al. Automated percutaneous lumbar discectomy versus chemonucleolysis in the treatment of sciatica. A randomized multicenter trial. *Spine (Phila Pa 1976).* Jan 1993;18(1):1-7. PMID 8434309
9. Freeman BJ, Mehdian R. Intradiscal electrothermal therapy, percutaneous discectomy, and nucleoplasty: what is the current evidence? *Curr Pain Headache Rep.* Jan 2008;12(1):14-21. PMID 18417018
10. Gibson JN, Waddell G. Surgical interventions for lumbar disc prolapse. *Cochrane Database Syst Rev.* 2007(2):CD001350. PMID 17443505
11. Boswell MV, Trescot AM, Datta S, et al. Interventional techniques: evidence-based practice guidelines in the management of chronic spinal pain. *Pain Physician.* Jan 2007;10(1):7-111. PMID 17256025
12. Haines SJ, Jordan N, Boen JR, et al. Discectomy strategies for lumbar disc herniation: results of the LAPDOG trial. *J Clin Neurosci.* Jul 2002;9(4):411-417. PMID 12217670
13. Chatterjee S, Foy PM, Findlay GF. Report of a controlled clinical trial comparing automated percutaneous lumbar discectomy and microdiscectomy in the treatment of contained lumbar disc herniation. *Spine (Phila Pa 1976).* Mar 15 1995;20(6):734-738. PMID 7604351
14. Manchikanti L, Abdi S, Atluri S, et al. An update of comprehensive evidence-based guidelines for interventional techniques in chronic spinal pain. Part II: guidance and recommendations. *Pain Physician.* Apr 2013; 16(2 Suppl):S49-283. PMID 23615883
15. Phan K, Xu J, Schultz K, et al. Full-endoscopic versus micro-endoscopic and open discectomy: A systematic review and meta-analysis of outcomes and complications. *Clin Neurol Neurosurg.* Mar 2017;154:1-12. PMID 28086154
16. Li XC, Zhong CF, Deng GB, et al. Full-endoscopic procedures versus traditional discectomy surgery for discectomy: a systematic review and meta-analysis of current global clinical trials. *Pain Physician.* Mar 2016; 19(3):103-118. PMID 27008284
17. Cong L, Zhu Y, Tu G. A meta-analysis of endoscopic discectomy versus open discectomy for symptomatic lumbar disk herniation. *Eur Spine J.* Jan 2016;25(1):134-143. PMID 25632840
18. Smith N, Masters J, Jensen C, et al. Systematic review of microendoscopic discectomy for lumbar disc herniation. *Eur Spine J.* Nov 2013;22(11):2458-2465. PMID 23793558
19. Teli M, Lovi A, Brayda-Bruno M, et al. Higher risk of dural tears and recurrent herniation with lumbar microendoscopic discectomy. *Eur Spine J.* Mar 2010;19(3):443-450. PMID 20127495
20. Garg B, Nagraja UB, Jayaswal A. Microendoscopic versus open discectomy for lumbar disc herniation: a prospective randomised study. *J Orthop Surg (Hong Kong).* Apr 2011;19(1):30-34. PMID 21519072

21. Gibson JN, Subramanian AS, Scott CE. A randomised controlled trial of transforaminal endoscopic discectomy vs. microdiscectomy. *Eur Spine J.* Mar 2017;26(3):847-856. PMID 27885470
22. Hussein M, Abdeldayem A, Mattar MM. Surgical technique and effectiveness of microendoscopic discectomy for large uncontained lumbar disc herniations: a prospective, randomized, controlled study with 8 years of follow-up. *Eur Spine J.* Sep 2014;23(9):1992-1999. PMID 24736930
23. Ruetten S, Komp M, Merk H, et al. Full-endoscopic cervical posterior foraminotomy for the operation of lateral disc herniations using 5.9-mm endoscopes: a prospective, randomized, controlled study. *Spine (Phila Pa 1976).* Apr 20 2008;33(9):940-948. PMID 18427313
24. Ruetten S, Komp M, Merk H, et al. Full-endoscopic interlaminar and transforaminal lumbar discectomy versus conventional microsurgical technique: a prospective, randomized, controlled study. *Spine (Phila Pa 1976).* Apr 20 2008;33(9):931-939. PMID 18427312
25. Ruetten S, Komp M, Merk H, et al. Recurrent lumbar disc herniation after conventional discectomy: a prospective, randomized study comparing full-endoscopic interlaminar and transforaminal versus microsurgical revision. *J Spinal Disord Tech.* Apr 2009;22(2):122-129. PMID 19342934
26. Ruetten S, Komp M, Merk H, et al. Full-endoscopic anterior decompression versus conventional anterior decompression and fusion in cervical disc herniations. *Int Orthop.* Dec 2009;33(6):1677-1682. PMID 19015851
27. Hermantin FU, Peters T, Quartararo L, et al. A prospective, randomized study comparing the results of open discectomy with those of video-assisted arthroscopic microdiscectomy. *J Bone Joint Surg Am.* Jul 1999; 81(7):958-965. PMID 10428127
28. Gotecha S, Ranade D, Patil SV, et al. The role of transforaminal percutaneous endoscopic discectomy in lumbar disc herniations. *J Craniovertebr Junction Spine.* 2016;7(4):217-223. PMID 27891030
29. Lee DY, Lee SH. Learning curve for percutaneous endoscopic lumbar discectomy. *Neurol Med Chir (Tokyo).* Sep 2008;48(9):383-388; discussion 388-389. PMID 18812679
30. Wang B, Lu G, Patel AA, et al. An evaluation of the learning curve for a complex surgical technique: the full endoscopic interlaminar approach for lumbar disc herniations. *Spine J.* Feb 2011;11(2):122-130. PMID 21296295
31. Tenenbaum S, Arzi H, Herman A, et al. Percutaneous posterolateral transforaminal endoscopic discectomy: clinical outcome, complications, and learning curve evaluation. *Surg Technol Int.* Dec 2011;21:278-283. PMID 22505002
32. Casal-Moro R, Castro-Menendez M, Hernandez-Blanco M, et al. Long-term outcome after microendoscopic discectomy for lumbar disk herniation: a prospective clinical study with a 5-year follow-up. *Neurosurgery.* Jun 2011;68(6):1568-1575; discussion 1575. PMID 21311384
33. Wang M, Zhou Y, Wang J, et al. A 10-year follow-up study on long-term clinical outcomes of lumbar microendoscopic discectomy. *J Neurol Surg A Cent Eur Neurosurg.* Aug 2012;73(4):195-198. PMID 22825836
34. Choi KC, Lee JH, Kim JS, et al. Unsuccessful percutaneous endoscopic lumbar discectomy: a single-center experience of 10,228 cases. *Neurosurgery.* Apr 2015;76(4):372-381. PMID 25599214
35. National Institute for Health and Care Excellence (NICE). Automated percutaneous mechanical lumbar discectomy-guidance [IPG141]. 2005; <http://guidance.nice.org.uk/IPG141/Guidance/pdf/English>. Accessed March 9, 2017.
36. National Institute for Health and Care Excellence (NICE). Percutaneous transforaminal endoscopic lumbar discectomy for sciatica [IPG556]. 2016; <https://www.nice.org.uk/guidance/ipg556>. Accessed March 9, 2017.
37. National Institute for Health and Care Excellence (NICE). Percutaneous interlaminar endoscopic lumbar discectomy for sciatica [IPG555]. 2016; <https://www.nice.org.uk/guidance/ipg555>. Accessed March 9, 2017.
38. Kreiner DS, Hwang SW, Easa JE, et al. An evidence-based clinical guideline for the diagnosis and treatment of lumbar disc herniation with radiculopathy. *Spine J.* Jan 2014;14(1):180-191. PMID 24239490

39. Chou R, Loeser JD, Owens DK, et al. Interventional therapies, surgery, and interdisciplinary rehabilitation for low back pain: an evidence-based clinical practice guideline from the American Pain Society. *Spine (Phila Pa 1976)*. May 1 2009;34(10):1066-1077. PMID 19363457
40. National Coverage Determination (NCD) for PERCUTANEOUS Image-Guided Lumbar Decompression for Lumbar Spinal Stenosis (150.13), Effective Date of this Version 12/7/2016.