

(80312)

Medical Benefit		Effective Date: 01/01/05	Next Review Date: 07/23
Preauthorization	No	Review Dates: 09/09, 09/10, 07/11, 07/12, 07/13, 07/14, 07/15, 07/16, 07/17, 07/18, 07/19, 07/20, 07/21, 07/22	

This protocol considers this test or procedure investigational. If the physician feels this service is medically necessary, preauthorization is recommended.

The following protocol contains medical necessity criteria that apply for this service. The criteria are also applicable to services provided in the local Medicare Advantage operating area for those members, unless separate Medicare Advantage criteria are indicated. If the criteria are not met, reimbursement will be denied and the patient cannot be billed. Please note that payment for covered services is subject to eligibility and the limitations noted in the patient's contract at the time the services are rendered.

RELATED PROTOCOL

None

Populations	Interventions	Comparators	Outcomes
Individuals: • With cerebral palsy	Interventions of interest are: • Hippotherapy	Comparators of interest are: • Standard clinical management	Relevant outcomes include: • Symptoms • Functional outcomes
Individuals: • With multiple sclerosis	Interventions of interest are: • Hippotherapy	Comparators of interest are: • Standard clinical management	Relevant outcomes include: • Symptoms • Functional outcomes
Individuals: • With stroke	Interventions of interest are: • Hippotherapy	Comparators of interest are: • Standard clinical management	Relevant outcomes include: • Symptoms • Functional outcomes
Individuals: • With gait and balance disorders other than cerebral palsy, multiple sclerosis, and stroke	Interventions of interest are: • Hippotherapy	Comparators of interest are: • Standard clinical management	Relevant outcomes include: • Symptoms • Functional outcomes

DESCRIPTION

Hippotherapy, also referred to as equine-assisted therapy, describes a treatment strategy that uses equine movement to engage sensory, neuromotor, and cognitive systems to achieve functional outcomes. Hippotherapy has been proposed as a therapy for patients with impaired walking or balance.

SUMMARY OF EVIDENCE

For individuals who have cerebral palsy, multiple sclerosis, stroke, or gait and balance disorders other than cerebral palsy, multiple sclerosis, and stroke who receive hippotherapy, the evidence includes systematic reviews,

randomized trials, and case series. Relevant outcomes include symptoms and functional outcomes. Studies in cerebral palsy, multiple sclerosis, stroke, and other indications have had variable findings. The randomized trials are generally small and have significant methodologic problems. In the largest randomized trial conducted to date (72 children), which had blinding outcome assessment, hippotherapy had no clinically significant impact on children with cerebral palsy. There are no randomized controlled trials showing that hippotherapy is superior to alternative treatments for patients with multiple sclerosis. Hippotherapy for other indications has been compared primarily with no intervention and, although some benefits have been seen, it has not been shown to be more effective than other active therapies. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

POLICY

Hippotherapy is considered **investigational**.

BACKGROUND

HIPPOTHERAPY

Hippotherapy has been proposed as a technique to decrease the energy requirements and improve walking in patients with cerebral palsy. It is thought that the natural swaying motion of the horse induces a pelvic movement in the rider that simulates human ambulation. Also, variations in the horse's movements can prompt natural equilibrium movements in the rider. Hippotherapy is also being evaluated in patients with multiple sclerosis and other causes of gait disorders, such as strokes.

As a therapeutic intervention, hippotherapy is typically conducted by a physical or occupational therapist and is aimed at improving impaired body function. Therapeutic horseback riding is typically conducted by riding instructors and is more frequently intended as social therapy. It is hoped that the multisensory environment may benefit children with profound social and communication deficits, such as autism spectrum disorder and schizophrenia. When considered together, hippotherapy and therapeutic riding are described as equine-assisted activities and therapies.

This protocol addresses equine-assisted activities that focus on improving physical functions such as balance and gait.

Services that are the subject of a clinical trial do not meet our Technology Assessment and Medically Necessary Services Protocol criteria and are considered investigational. *For explanation of experimental and investigational, please refer to the Technology Assessment and Medically Necessary Services Protocol.*

It is expected that only appropriate and medically necessary services will be rendered. We reserve the right to conduct prepayment and postpayment reviews to assess the medical appropriateness of the above-referenced procedures. **Some of this protocol may not pertain to the patients you provide care to, as it may relate to products that are not available in your geographic area.**

REFERENCES

We are not responsible for the continuing viability of web site addresses that may be listed in any references below.

1. Selph SS, Skelly AC, Wasson N, et al. Physical Activity and the Health of Wheelchair Users: A Systematic Review in Multiple Sclerosis, Cerebral Palsy, and Spinal Cord Injury. Number 241. Agency for Healthcare Research and Quality, US Department of Health and Human Services; 2021. Accessed February 16, 2022. <https://www.ncbi.nlm.nih.gov/books/NBK574875/#ch4.s5>
2. Tseng SH, Chen HC, Tam KW. Systematic review and meta-analysis of the effect of equine assisted activities and therapies on gross motor outcome in children with cerebral palsy. *Disabil Rehabil.* Jan 2013;35(2):89-99. PMID 22630812
3. Zadnikar M, Kastrin A. Effects of hippotherapy and therapeutic horseback riding on postural control or balance in children with cerebral palsy: a meta-analysis. *Dev Med Child Neurol.* Aug 2011;53(8):684-91. PMID 21729249
4. Chinniah H, Natarajan M, Ramanathan R, et al. Effects of horse riding simulator on sitting motor function in children with spastic cerebral palsy. *Physiother Res Int.* Oct 2020;25(4):e1870. PMID 32808394
5. Kwon JY, Chang HJ, Yi SH, et al. Effect of hippotherapy on gross motor function in children with cerebral palsy: a randomized controlled trial. *J Altern Complement Med.* Jan 2015;21(1):15-21. PMID 25551626
6. Davis E, Davies B, Wolfe R, et al. A randomized controlled trial of the impact of therapeutic horse riding on the quality of life, health, and function of children with cerebral palsy. *Dev Med Child Neurol.* Feb 2009; 51(2):111-9; discussion 88. PMID 19191844
7. McGibbon NH, Benda W, Duncan BR, et al. Immediate and long-term effects of hippotherapy on symmetry of adductor muscle activity and functional ability in children with spastic cerebral palsy. *Arch Phys Med Rehabil.* Jun 2009;90(6):966-74. PMID 19480872
8. Bronson C, Brewerton K, Ong J, et al. Does hippotherapy improve balance in persons with multiple sclerosis: a systematic review. *Eur J Phys Rehabil Med.* Sep 2010;46(3):347-53. PMID 20927000
9. Silkwood-Sherer D, Warmbier H. Effects of hippotherapy on postural stability, in persons with multiple sclerosis: a pilot study. *J Neurol Phys Ther.* Jun 2007;31(2):77-84. PMID 17558361
10. Munoz-Lasa S, Ferriero G, Valero R, et al. Effect of therapeutic horseback riding on balance and gait of people with multiple sclerosis. *G Ital Med Lav Ergon.* Oct-Dec 2011;33(4):462-7. PMID 22452106
11. Frevel D, Maurer M. Internet-based home training is capable to improve balance in multiple sclerosis: a randomized controlled trial. *Eur J Phys Rehabil Med.* Feb 2015;51(1):23-30. PMID 24755773
12. Lee CW, Kim SG, Yong MS. Effects of hippotherapy on recovery of gait and balance ability in patients with stroke. *J Phys Ther Sci.* Feb 2014;26(2):309-11. PMID 24648655
13. Bunketorp-Kall L, Pekna M, Pekny M, et al. Effects of horse-riding therapy and rhythm and music-based therapy on functional mobility in late phase after stroke. *NeuroRehabilitation.* Dec 18 2019;45(4):483-492. PMID 31868694
14. Bunketorp Kall L, Lundgren-Nilsson A, Blomstrand C, et al. The effects of a rhythm and music-based therapy program and therapeutic riding in late recovery phase following stroke: a study protocol for a three-armed randomized controlled trial. *BMC Neurol.* Nov 21 2012;12:141. PMID 23171380
15. Wood WH, Fields BE. Hippotherapy: a systematic mapping review of peer-reviewed research, 1980 to 2018. *Disabil Rehabil.* May 2021;43(10):1463-1487. PMID 31491353
16. Kim SG, Lee CW. The effects of hippotherapy on elderly persons' static balance and gait. *J Phys Ther Sci.* Jan 2014;26(1):25-7. PMID 24567669
17. Araujo TB, Silva NA, Costa JN, et al. Effect of equine-assisted therapy on the postural balance of the elderly. *Rev Bras Fisioter.* Sep-Oct 2011;15(5):414-9. PMID 22002189
18. de Araujo TB, de Oliveira RJ, Martins WR, et al. Effects of hippotherapy on mobility, strength and balance in elderly. *Arch Gerontol Geriatr.* May-Jun 2013;56(3):478-81. PMID 23290005
19. Homnick DN, Henning KM, Swain CV, et al. Effect of therapeutic horseback riding on balance in community-dwelling older adults with balance deficits. *J Altern Complement Med.* Jul 2013;19(7):622-6. PMID 23360659
20. Abdel-Aziem AA, Abdelraouf OR, Ghally SA, et al. A 10-Week Program of Combined Hippotherapy and Scroth's Exercises Improves Balance and Postural Asymmetries in Adolescence Idiopathic Scoliosis: A Randomized Controlled Study. *Children (Basel).* Dec 30 2021;9(1). PMID 35053648

21. Silkwood-Sherer DJ, Killian CB, Long TM, et al. Hippotherapy--an intervention to habilitate balance deficits in children with movement disorders: a clinical trial. *Phys Ther.* May 2012;92(5):707-17. PMID 22247403
22. Giagazoglou P, Arabatzi F, Dipla K, et al. Effect of a hippotherapy intervention program on static balance and strength in adolescents with intellectual disabilities. *Res Dev Disabil.* Nov-Dec 2012;33(6):2265-70. PMID 22853887
23. Lechner HE, Kakebeeke TH, Hegemann D, et al. The effect of hippotherapy on spasticity and on mental well-being of persons with spinal cord injury. *Arch Phys Med Rehabil.* Oct 2007;88(10):1241-8. PMID 17908564
24. American Hippotherapy Association, Inc. Statements of best practice for the use of hippotherapy by occupational therapy, physical therapy, and speech-language pathology professionals. Revised February 2021. <https://www.americanhippotherapyassociation.org/assets/docs/AHA%20Statements%20of%20Best%20Practice%20February%202021.pdf>. Accessed January 31, 2022.