

(10405)

Medical Benefit		Effective Date: 08/01/20	Next Review Date: 05/23
Preauthorization	No	Review Dates: 09/09, 05/10, 05/11, 05/12, 05/13, 05/14, 05/15, 05/16, 05/17, 05/18, 05/19, 05/20, 05/21, 05/22	

Preauthorization is not required.

The following protocol contains medical necessity criteria that apply for this service. The criteria are also applicable to services provided in the local Medicare Advantage operating area for those members, unless separate Medicare Advantage criteria are indicated. If the criteria are not met, reimbursement will be denied and the patient cannot be billed. Please note that payment for covered services is subject to eligibility and the limitations noted in the patient's contract at the time the services are rendered.

RELATED PROTOCOLS

Functional Neuromuscular Electrical Stimulation

Myoelectric Prosthetic and Orthotic Components for the Upper Limb

Populations	Interventions	Comparators	Outcomes
Individuals: • With transfemoral amputation	Interventions of interest are: • Prosthesis with a microprocessor-controlled knee	Comparators of interest are: • Prosthesis with a conventional knee	Relevant outcomes include: • Functional outcomes • Health status measures • Quality of life
Individuals: • With transfemoral amputation	Interventions of interest are: • Prosthesis with a powered knee	Comparators of interest are: • Prosthesis with a conventional knee	Relevant outcomes include: • Functional outcomes • Health status measures • Quality of life
Individuals: • With tibial amputation	Interventions of interest are: • Prosthesis with a microprocessor-controlled ankle-foot	Comparators of interest are: • Prosthesis with a conventional foot-ankle	Relevant outcomes include: • Functional outcomes • Health status measures • Quality of life
Individuals: • With tibial amputation	Interventions of interest are: • Prosthesis with a powered ankle-foot	Comparators of interest are: • Prosthesis with a conventional ankle-foot	Relevant outcomes include: • Functional outcomes • Health status measures • Quality of life

DESCRIPTION

Microprocessor-controlled prostheses use feedback from sensors to adjust joint movement on a real-time as-needed basis. Active joint control is intended to improve safety and function, particularly for patients who can maneuver on uneven terrain and with variable gait.

SUMMARY OF EVIDENCE

For individuals who have a transfemoral amputation who receive a prosthesis with a microprocessor-controlled knee, the evidence includes a number of within-subject comparisons of microprocessor-controlled knees versus non-microprocessor-controlled knee joints. Relevant outcomes are functional outcomes, health status measures, and quality of life. For K3- and K4-level amputees, studies have shown an objective improvement in function on some outcome measures, particularly for hill and ramp descent, and strong patient preference for microprocessor-controlled prosthetic knees. Benefits include a more normal gait, increased stability, and a decrease in falls. The evidence in Medicare level K2 ambulators suggests that a prosthesis with stance control only can improve activities that require balance and improve walking in this population. For these reasons, a microprocessor-controlled knee may provide incremental benefit for these individuals. The potential to achieve a higher functional level with a microprocessor-controlled knee includes having the appropriate physical and cognitive ability to use the advanced technology. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have a transfemoral amputation who receive a prosthesis with a powered knee, the evidence includes no data. Relevant outcomes are functional outcomes, health status measures, and quality of life. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have a tibial amputation who receive a prosthesis with a microprocessor-controlled ankle-foot, the evidence includes limited data. Relevant outcomes are functional outcomes, health status measures, and quality of life. The limited evidence available to date does not support an improvement in functional outcomes using microprocessor-controlled ankle-foot prostheses compared with standard prostheses. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have a tibial amputation who receive a prosthesis with a powered ankle-foot, the evidence includes limited data. Relevant outcomes are functional outcomes, health status measures, and quality of life. The limited evidence available to date does not support an improvement in functional outcomes using powered ankle-foot prostheses compared with standard prostheses. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

POLICY

A microprocessor-controlled knee may be considered **medically necessary** in individuals with transfemoral amputation who meet the following requirements:

- demonstrated need for long distance ambulation at variable rates (use of the limb in the home or for basic community ambulation is not sufficient to justify provision of the computerized limb over standard limb applications) OR demonstrated patient need for regular ambulation on uneven terrain or for regular use on stairs (use of the limb for limited stair climbing in the home or employment environment is not sufficient evidence for prescription of this device over standard prosthetic application); AND
- physical ability, including adequate cardiovascular and pulmonary reserve, for ambulation at faster than normal walking speed; AND
- adequate cognitive ability to master use and care requirements for the technology.

A microprocessor-controlled knee is considered **not medically necessary** in individuals who do not meet these criteria.

A powered knee is considered **investigational**.

A microprocessor-controlled or powered ankle-foot is considered **investigational**.

POLICY GUIDELINES

Amputees should be evaluated by an independent qualified professional to determine the most appropriate prosthetic components and control mechanism. A trial period may be indicated to evaluate the tolerability and efficacy of the prosthesis in a real-life setting. Decisions about the potential benefits of microprocessor-knees involve multiple factors including activity levels as well as the patient's physical and cognitive ability. A patient's need for daily ambulation of at least 400 continuous yards, daily and frequent ambulation at variable cadence or on uneven terrain (e.g., gravel, grass, curbs), and daily and frequent use of ramps and/or stairs (especially stair descent) should be considered as part of the decision. Typically, daily and frequent need of two or more of these activities would be needed to show benefit.

PATIENT SELECTION AND IDENTIFICATION

For patients in whom the potential benefits of the microprocessor knees are uncertain, patients may first be fitted with a standard prosthesis to determine their level of function with the standard device.

The following are guidelines from the Veteran's Health Administration Prosthetic Clinical Management Program Clinical Practice Recommendations for Microprocessor Knees. (Berry, 2006).

- A. Contraindications for use of the microprocessor knee should include the following:
- a. Any condition that prevents socket fitting, such as a complicated wound or intractable pain which precludes socket wear.
 - b. Inability to tolerate the weight of the prosthesis.
 - c. Medicare Level K0—no ability or potential to ambulate or transfer.
 - d. Medicare Level K1—limited ability to transfer or ambulate on level ground at fixed cadence.
 - e. Medicare Level K2—limited community ambulator that does not have the cardiovascular reserve, strength, and balance to improve stability in stance to permit increased independence, less risk of falls, and potential to advance to a less-restrictive walking device.
 - f. Inability to use swing and stance features of the knee unit.
 - g. Poor balance or ataxia that limits ambulation.
 - h. Significant hip flexion contracture (over 20°).
 - i. Significant deformity of remaining limb that would impair ability to stride.
 - j. Limited cardiovascular and/or pulmonary reserve or profound weakness.
 - k. Limited cognitive ability to understand gait sequencing or care requirements.
 - l. Long distance or competitive running.
 - m. Falls outside of recommended weight or height guidelines of manufacturer.
 - n. Specific environmental factors—such as excessive moisture or dust, or inability to charge the prosthesis.
 - o. Extremely rural conditions where maintenance ability is limited.
- B. Indications for use of the microprocessor knee should include the following:
- a. Adequate cardiovascular and pulmonary reserve to ambulate at variable cadence.
 - b. Adequate strength and balance in stride to activate the knee unit.
 - c. Should not exceed the weight or height restrictions of the device.

- d. Adequate cognitive ability to master technology and gait requirements of device.
 - e. Hemi-pelvectomy through knee-disarticulation level of amputation, including bilateral; lower extremity amputees are candidates if they meet functional criteria as listed.
 - f. Patient is an active walker and requires a device that reduces energy consumption to permit longer distances with less fatigue.
 - g. Daily activities or job tasks that do not permit full focus of concentration on knee control and stability—such as uneven terrain, ramps, curbs, stairs, repetitive lifting, and/or carrying.
 - h. Medicare Level K2—limited community ambulator, but only if improved stability in stance permits increased independence, less risk of falls, and potential to advance to a less restrictive walking device, and patient has cardiovascular reserve, strength, and balance to use the prosthesis. *The microprocessor enables fine-tuning and adjustment of the hydraulic mechanism to accommodate the unique motor skills and demands of the functional level K2 ambulator.*
 - i. Medicare Level K3—unlimited community ambulator.
 - j. Medicare Level K4—active adult, athlete, who has the need to function as a K3 level in daily activities.
 - k. Potential to lessen back pain by providing more secure stance control, using less muscle control to keep knee stable.
 - l. Potential to unload and decrease stress on remaining limb.
 - m. Potential to return to an active lifestyle.
- C. Physical and Functional Fitting Criteria for New Amputees:
- a. New amputees may be considered if they meet certain criteria as outlined above.
 - b. Premorbid and current functional assessment important determinant.
 - c. Requires stable wound and ability to fit socket.
 - d. Immediate postoperative fit is possible.
 - e. Must have potential to return to active lifestyle.

MEDICARE ADVANTAGE

For Medicare Advantage a lower limb prosthesis would only be considered **medically necessary** when the member:

1. Will reach or maintain a defined functional state within a reasonable period of time; and
2. Is motivated to ambulate.

KNEE

A fluid, pneumatic, or electronic/microprocessor knee may be **medically necessary** for patients whose functional level is three or above (see Medicare Advantage Policy Guidelines).

The powered and programmable flexion/extension assist control addition is **medically necessary** only when the patient meets all of the criteria below:

- Has a microprocessor (swing and stance phase type controlled electronic) knee
- K3 functional level only

- Weight greater than 110 lbs. and less than 275 lbs.
- Has a documented comorbidity of the spine and/or sound limb affecting hip extension and/or quadriceps function that impairs K3 level function with the use of a microprocessor-controlled knee alone
- Is able to make use of a product that requires daily charging
- Is able to understand and respond to error alerts and alarms indicating problems with the function of the unit

FOOT

A microprocessor controlled ankle foot system may be **medically necessary** for patients whose functional level is three or above (see Medicare Advantage Policy Guidelines).

MEDICARE ADVANTAGE POLICY GUIDELINES

Potential functional ability is based on the reasonable expectations of the prosthetist, and treating physician, considering factors including, but not limited to:

1. The member's past history (including prior prosthetic use if applicable); and
2. The member's current condition including the status of the residual limb and the nature of other medical problems; and
3. The member's desire to ambulate.

REHABILITATION CLASSIFICATION LEVELS

Level 0: Does not have the ability or potential to ambulate or transfer safely with or without assistance and prosthesis does not enhance their quality of life or mobility.

Level 1: Has the ability or potential to use prosthesis for transfers or ambulation on level surfaces at fixed cadence. Typical of the limited and unlimited household ambulator.

Level 2: Has the ability or potential for ambulation with the ability to traverse low level environmental barriers such as curbs, stairs or uneven surfaces. Typical of the limited community ambulator.

Level 3: Has the ability or potential for ambulation with variable cadence. Typical of the community ambulator who has the ability to traverse most environmental barriers and may have vocational, therapeutic, or exercise activity that demands prosthetic utilization beyond simple locomotion.

Level 4: Has the ability or potential for prosthetic ambulation that exceeds basic ambulation skills, exhibiting high impact, stress, or energy levels. Typical of the prosthetic demands of the child, active adult, or athlete.

BACKGROUND

LOWER-EXTREMITY PROSTHETICS

More than 100 different prosthetic ankle-foot and knee designs are currently available. The choice of the most appropriate design may depend on the patient's underlying activity level. For example, the requirements of a prosthetic knee in an elderly, largely homebound individual will differ from those of a younger, active person. Key elements of prosthetic knee design involve providing stability during both the stance and swing phase of the gait. Prosthetic knees vary in their ability to alter the cadence of the gait, or the ability to walk on rough or uneven surfaces. In contrast to more simple prostheses, which are designed to function optimally at 1 walking cadence, fluid and hydraulic-controlled devices are designed to allow amputees to vary their walking speed by

matching the movement of the shin portion of the prosthesis to the movement of the upper leg. For example, the rate at which the knee flexes after “toe-off” and then extends before heel strike depends in part on the mechanical characteristics of the prosthetic knee joint. If the resistance to flexion and extension of the joint does not vary with gait speed, the prosthetic knee extends too quickly or too slowly relative to the heel strike if the cadence is altered. When properly controlled, hydraulic or pneumatic swing-phase controls allow the prosthetist to set a pace adjusted to the individual amputee, from very slow to a race-walking pace. Hydraulic prostheses are heavier than other options and require gait training; for these reasons, these prostheses are prescribed for athletic or fit individuals. Other design features include multiple centers of rotation, referred to as “polycentric knees.” The mechanical complexity of these devices allows engineers to optimize selected stance and swing-phase features.

REGULATORY STATUS

According to the manufacturers, microprocessor-controlled prostheses are considered a class I device by the FDA and are exempt from 510(k) requirements. This classification does not require submission of clinical data regarding efficacy but only notification of FDA prior to marketing. FDA product codes: ISW, KFX.

Services that are the subject of a clinical trial do not meet our Technology Assessment and Medically Necessary Services Protocol criteria and are considered investigational. *For explanation of experimental and investigational, please refer to the Technology Assessment and Medically Necessary Services Protocol.*

It is expected that only appropriate and medically necessary services will be rendered. We reserve the right to conduct prepayment and postpayment reviews to assess the medical appropriateness of the above-referenced procedures. **Some of this protocol may not pertain to the patients you provide care to, as it may relate to products that are not available in your geographic area.**

REFERENCES

We are not responsible for the continuing viability of web site addresses that may be listed in any references below.

1. Flynn K. Short Report: Computerized lower limb prosthesis (VA Technology Assessment Program). No. 2. Boston, MA: Veterans Health Administration; 2000.
2. Theeven P, Hemmen B, Rings F, et al. Functional added value of microprocessor-controlled knee joints in daily life performance of Medicare Functional Classification Level-2 amputees. *J Rehabil Med*. Oct 2011; 43(10):906-15. PMID 21947182
3. Theeven PJ, Hemmen B, Geers RP, et al. Influence of advanced prosthetic knee joints on perceived performance and everyday life activity level of low-functional persons with a transfemoral amputation or knee disarticulation. *J Rehabil Med*. May 2012;44(5):454-61. PMID 22549656
4. Burnfield JM, Eberly VJ, Gronely JK, et al. Impact of stance phase microprocessor-controlled knee prosthesis on ramp negotiation and community walking function in K2 level transfemoral amputees. *Prosthet Orthot Int*. Mar 2012;36(1):95-104. PMID 2223685
5. Orendurff MS, Segal AD, Klute GK, et al. Gait efficiency using the C-Leg. *J Rehabil Res Dev*. Mar-Apr 2006; 43(2):239-46. PMID 16847790
6. Klute GK, Berge JS, Orendurff MS, et al. Prosthetic intervention effects on activity of lower-extremity amputees. *Arch Phys Med Rehabil*. May 2006;87(5):717-22. PMID 16635636

7. Williams RM, Turner AP, Orendurff M, et al. Does having a computerized prosthetic knee influence cognitive performance during amputee walking?. *Arch Phys Med Rehabil*. Jul 2006;87(7):989-94. PMID 16813788
8. Hafner BJ, Smith DG. Differences in function and safety between Medicare Functional Classification Level-2 and -3 transfemoral amputees and influence of prosthetic knee joint control. *J Rehabil Res Dev*. 2009;46(3):417-33. PMID 19675993
9. Highsmith MJ, Kahle JT, Miro RM, et al. Ramp descent performance with the C-Leg and interrater reliability of the Hill Assessment Index. *Prosthet Orthot Int*. Oct 2013;37(5):362-8. PMID 23327837
10. Howard CL, Wallace C, Perry B, et al. Comparison of mobility and user satisfaction between a microprocessor knee and a standard prosthetic knee: a summary of seven single-subject trials. *Int J Rehabil Res*. Mar 2018;41(1):63-73. PMID 29293160
11. Hafner BJ, Willingham LL, Buell NC, et al. Evaluation of function, performance, and preference as transfemoral amputees transition from mechanical to microprocessor control of the prosthetic knee. *Arch Phys Med Rehabil*. Feb 2007;88(2):207-17. PMID 17270519
12. Kaufman KR, Bernhardt KA, Symms K. Functional assessment and satisfaction of transfemoral amputees with low mobility (FASTK2): A clinical trial of microprocessor-controlled vs. non-microprocessor-controlled knees. *Clin Biomech (Bristol, Avon)*. Oct 2018;58:116-122. PMID 30077128
13. Kaufman KR, Levine JA, Brey RH, et al. Gait and balance of transfemoral amputees using passive mechanical and microprocessor-controlled prosthetic knees. *Gait Posture*. Oct 2007;26(4):489-93. PMID 17869114
14. Kaufman KR, Levine JA, Brey RH, et al. Energy expenditure and activity of transfemoral amputees using mechanical and microprocessor-controlled prosthetic knees. *Arch Phys Med Rehabil*. Jul 2008;89(7):1380-5. PMID 18586142
15. Johansson JL, Sherrill DM, Riley PO, et al. A clinical comparison of variable-damping and mechanically passive prosthetic knee devices. *Am J Phys Med Rehabil*. Aug 2005;84(8):563-75. PMID 16034225
16. Carse B, Scott H, Brady L, et al. Evaluation of gait outcomes for individuals with established unilateral transfemoral amputation following the provision of microprocessor controlled knees in the context of a clinical service. *Prosthet Orthot Int*. Jun 01 2021;45(3):254-261. PMID 34016870
17. Hofstad C, Linde H, Limbeek J, et al. Prescription of prosthetic ankle-foot mechanisms after lower limb amputation. *Cochrane Database Syst Rev*. 2004;(1):CD003978. PMID 14974050
18. Alimusaj M, Fradet L, Braatz F, et al. Kinematics and kinetics with an adaptive ankle foot system during stair ambulation of transtibial amputees. *Gait Posture*. Oct 2009;30(3):356-63. PMID 19616436
19. Fradet L, Alimusaj M, Braatz F, et al. Biomechanical analysis of ramp ambulation of transtibial amputees with an adaptive ankle foot system. *Gait Posture*. Jun 2010;32(2):191-8. PMID 20457526
20. Darter BJ, Wilken JM. Energetic consequences of using a prosthesis with adaptive ankle motion during slope walking in persons with a transtibial amputation. *Prosthet Orthot Int*. Feb 2014;38(1):5-11. PMID 23525888
21. Gailey RS, Gaunard I, Agrawal V, et al. Application of self-report and performance-based outcome measures to determine functional differences between four categories of prosthetic feet. *J Rehabil Res Dev*. 2012;49(4):597-612. PMID 22773262
22. Delussu AS, Brunelli S, Paradisi F, et al. Assessment of the effects of carbon fiber and bionic foot during overground and treadmill walking in transtibial amputees. *Gait Posture*. Sep 2013;38(4):876-82. PMID 23702342
23. Thomas-Pohl M, Villa C, Davot J, et al. Microprocessor prosthetic ankles: comparative biomechanical evaluation of people with transtibial traumatic amputation during standing on level ground and slope. *Disabil Rehabil Assist Technol*. Jan 2021;16(1):17-26. PMID 31535903
24. Au S, Berniker M, Herr H. Powered ankle-foot prosthesis to assist level-ground and stair-descent gaits. *Neural Netw*. May 2008;21(4):654-66. PMID 18499394
25. Ferris AE, Aldridge JM, Rabago CA, et al. Evaluation of a powered ankle-foot prosthetic system during walking. *Arch Phys Med Rehabil*. Nov 2012;93(11):1911-8. PMID 22732369
26. Herr HM, Grabowski AM. Bionic ankle-foot prosthesis normalizes walking gait for persons with leg amputation. *Proc Biol Sci*. Feb 07 2012;279(1728):457-64. PMID 21752817

27. Mancinelli C, Patritti BL, Tropea P, et al. Comparing a passive-elastic and a powered prosthesis in transtibial amputees. *Annu Int Conf IEEE Eng Med Biol Soc.* 2011;2011:8255-8. PMID 22256259
28. Webster JB, Crunkhorn A, Sall J, et al. Clinical Practice Guidelines for the Rehabilitation of Lower Limb Amputation: An Update from the Department of Veterans Affairs and Department of Defense. *Am J Phys Med Rehabil.* Sep 2019;98(9):820-829. PMID 31419214
29. Noridian Healthcare Solutions, LLC, (Jurisdiction A - New York - Entire State, Connecticut, Maine, Massachusetts, New Hampshire, Rhode Island, Vermont, District of Columbia, Delaware, Maryland, New Jersey, Pennsylvania) (Jurisdiction D - Alaska, American Samoa, Arizona, California - Entire State, Guam, Hawaii, Iowa, Idaho, Kansas, Missouri - entire state, Montana, North Dakota, Nebraska, Nevada, Oregon, South Dakota, Utah, Washington, Wyoming, Northern Mariana Islands) Local Coverage Determination (LCD): Lower LIMB Prostheses (L33787), Revision Effective Date for services performed on or after 01/01/2020.