

Protocol

Percutaneous Vertebroplasty and Sacroplasty

(60125)

Medical Benefit		Effective Date: 10/01/20	Next Review Date: 07/21
Preauthorization	No	Review Dates: 04/07, 05/08, 01/09, 01/10, 09/10, 07/11, 07/12, 07/13, 07/14, 07/15, 07/16, 07/17, 07/18, 07/19, 07/20	

Preauthorization is not required.

The following protocol contains medical necessity criteria that apply for this service. The criteria are also applicable to services provided in the local Medicare Advantage operating area for those members, unless separate Medicare Advantage criteria are indicated. If the criteria are not met, reimbursement will be denied and the patient cannot be billed. Please note that payment for covered services is subject to eligibility and the limitations noted in the patient's contract at the time the services are rendered.

RELATED PROTOCOLS

Diagnosis and Treatment of Sacroiliac Joint Pain

Percutaneous Balloon Kyphoplasty, Radiofrequency Kyphoplasty, and Mechanical Vertebral Augmentation

Populations	Interventions	Comparators	Outcomes
Individuals: • With symptomatic osteoporotic vertebral fractures between six weeks and one year old	Interventions of interest are: • Vertebroplasty	Comparators of interest are: • Conservative management	Relevant outcomes include: • Symptoms • Functional outcomes • Quality of life • Hospitalizations • Medication use • Treatment-related morbidity
Individuals: • With symptomatic osteoporotic vertebral fractures less than six weeks old	Interventions of interest are: • Vertebroplasty	Comparators of interest are: • Conservative management	Relevant outcomes include: • Symptoms • Functional outcomes • Quality of life • Hospitalizations • Medication use • Treatment-related morbidity
Individuals: • With sacral insufficiency fractures	Interventions of interest are: • Sacroplasty	Comparators of interest are: • Conservative management	Relevant outcomes include: • Symptoms • Functional outcomes • Quality of life • Hospitalizations • Medication use • Treatment-related morbidity

DESCRIPTION

Percutaneous vertebroplasty is an interventional technique involving the fluoroscopically guided injection of

polymethyl methacrylate into a weakened vertebral body. The technique has been investigated to provide mechanical support and symptomatic relief in patients with osteoporotic vertebral compression fractures or those with osteolytic lesions of the spine (e.g., multiple myeloma, metastatic malignancies); as a treatment for sacral insufficiency fractures; and as a technique to limit blood loss related to surgery.

SUMMARY OF EVIDENCE

For individuals who have symptomatic osteoporotic vertebral fractures between six weeks and one year old who receive vertebroplasty, the evidence includes two randomized sham-controlled trials, nonblinded randomized controlled trials (RCTs) comparing vertebroplasty with conservative management, and systematic reviews of these RCTs. Relevant outcomes are symptoms, functional outcomes, quality of life, hospitalizations, medication use, and treatment-related morbidity. Despite the completion of multiple RCTs, including two with sham controls, the efficacy of vertebroplasty for painful osteoporotic compression fractures remains uncertain. Two meta-analysis studies which included the two sham-controlled trials have demonstrated mixed results. The two studies had methodologic issues, including the choice of sham procedure and the potential of the sham procedure to have a therapeutic effect by reducing pain. Questions have also been raised about the low percentage of patients screened who participated in the trial, the volume of polymethylmethacrylate injected, and the inclusion of patients with chronic pain. Overall, conclusions about the effect of vertebroplasty remain unclear. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals with symptomatic osteoporotic vertebral fractures less than six weeks old who receive vertebroplasty, the evidence includes a randomized sham-controlled trial and nonblinded RCTs comparing vertebroplasty with conservative management. Relevant outcomes are symptoms, functional outcomes, quality of life, hospitalizations, medication use, and treatment-related morbidity. For acute fractures, conservative therapy consisting of rest, analgesics, and physical therapy is an option, and symptoms will resolve in a large percentage of patients with conservative treatment only. However, a sham-controlled randomized trial in patients who had severe pain of fewer than six weeks in duration found a significant benefit of vertebroplasty for the treatment of osteoporotic vertebral fracture at the thoracolumbar junction. Other RCTs without sham controls have reported that vertebroplasty is associated with significant improvements in pain and reductions in the duration of bed rest. Given the high morbidity associated with extended bed rest in older adults, this procedure is considered to have a significant health benefit. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals with sacral insufficiency fractures who receive sacroplasty, the evidence includes two prospective cohort studies and a case series. Relevant outcomes are symptoms, functional outcomes, quality of life, hospitalizations, medication use, and treatment-related morbidity. No RCTs have been reported. The available evidence includes a prospective cohort study and a retrospective series of 243 patients. These studies have reported rapid and sustained decreases in pain following percutaneous sacroplasty. Additional literature has mostly reported immediate improvements following the procedure. However, due to the small size of the evidence base, the harms associated with sacroplasty have not been adequately studied. The evidence is insufficient to determine the effects of the technology on health outcomes.

Vertebroplasty has been investigated as an intervention to provide mechanical support and symptomatic relief in patients with an osteoporotic vertebral compression fracture and in those with osteolytic lesions of the spine (i.e., multiple myeloma, metastatic malignancies). Clinical input obtained in 2008 provided uniform support for the use of vertebroplasty in painful osteoporotic fractures. Reconsideration of the available evidence and evaluation of the input led to a conclusion that, consistent results of numerous case series, including large prospective reports, the evidence was sufficient to determine that vertebroplasty is a reasonable treatment option in patients with vertebral fractures who have failed to respond to conservative treatment (at least six weeks with analgesics, physical therapy, and rest). It is also clinically reasonable to consider the evidence supporting the

clinical benefit of vertebroplasty in the osteoporotic vertebral fracture to support its use in osteolytic lesions of the spine (e.g., multiple myeloma, metastatic malignancies).

POLICY

Percutaneous vertebroplasty may be considered **medically necessary** for the treatment of symptomatic osteoporotic vertebral fractures that have failed to respond to conservative treatment (e.g., analgesics, physical therapy; rest) for at least six weeks.

Percutaneous vertebroplasty may be considered **medically necessary** for the treatment of symptomatic osteoporotic vertebral fractures that are less than six weeks in duration that have led to hospitalization or persist at a level that prevents ambulation.

Percutaneous vertebroplasty may be considered **medically necessary** for the treatment of severe pain due to osteolytic lesions of the spine related to multiple myeloma or metastatic malignancies.

Percutaneous vertebroplasty is considered **investigational** for all other indications, including use in acute vertebral fractures due to osteoporosis or trauma.

Percutaneous sacroplasty is considered **investigational** for all indications, including use in sacral insufficiency fractures due to osteoporosis and sacral lesions due to metastatic malignancies or multiple myeloma.

MEDICARE ADVANTAGE

The following only addresses Percutaneous Vertebral Augmentation (PVA) for Osteoporotic Vertebral Compression Fracture (VCF). See above for other indications.

PVA [percutaneous vertebroplasty (PVP)] is considered **medically necessary** with BOTH the following:

1. Inclusion criteria (ALL are required):
 - a. Acute* (less than six weeks) osteoporotic Vertebral Compression Fracture (VCF) (T5 – L5) by recent (within 30 days) advanced imaging (bone marrow edema on MRI or bone-scan/SPECT/CT uptake)
 - b. Symptomatic (ONE):
 - i. Hospitalized with severe pain (Numeric Rating Scale (NRS) or Visual Analog Scale (VAS) pain score ≥ 8)
 - ii. Non-hospitalized with moderate to severe pain (NRS or VAS ≥ 5) despite optimal non-surgical management** (ONE):
 1. Worsening pain
 2. Stable to improved pain (but NRS or VAS still ≥ 5) (with two or more of the following):
 - A. Progression of vertebral body height loss
 - B. More than 25% vertebral body height reduction
 - C. Kyphotic deformity
 - D. Severe impact of VCF on daily functioning (Roland Morris Disability Questionnaire (RDQ)) >17
 - c. Multidisciplinary team consensus (ALL are required)
 - i. Referring physician (e.g., rheumatologist, endocrinologist)

- ii. Treating physician (i.e., performing the PVA)
 - iii. Radiologist
 - iv. Neurologist
2. Exclusion criteria (Can have NONE of the following):
- a. Absolute contraindication
 - i. Current back pain is not primarily due to the identified acute VCF(s).
 - ii. Osteomyelitis, discitis or active systemic infection
 - iii. Pregnancy
 - iv. Greater than three vertebral fractures
 - b. Relative contraindication
 - i. Allergy to bone cement or opacification agents
 - ii. Coagulopathy
 - iii. Spinal instability
 - iv. Myelopathy from the fracture
 - v. Neurologic deficit
 - vi. Neural impingement
 - vii. Fracture retropulsion/canal compromise

*at least an acute component (e.g., acute on chronic)

**consider including pedicle periosteal infiltration

Percutaneous sacroplasty is considered **not medically necessary**.

BACKGROUND

TREATMENT OF VERTEBRAL COMPRESSION FRACTURE

Chronic symptoms do not tend to respond to the management strategies for acute pain such as bed rest, immobilization or bracing device, and analgesic medication, sometimes including narcotic analgesics. The source of chronic pain after vertebral compression fracture may not be from the vertebra itself but may be predominantly related to strain on muscles and ligaments secondary to kyphosis. This type of pain frequently does not improve with analgesics and may be better addressed through exercise or physical therapy. Improvements in pain and ability to function are the principal outcomes of interest for treatment of osteoporotic fractures.

TREATMENT OF SACRAL INSUFFICIENCY FRACTURES

Similar interventions are used for sacral and vertebral fractures and include bed rest, bracing, and analgesics. Initial clinical improvements may occur quickly; however, resolution of all symptoms may not occur for nine to 12 months.^{1,2}

VERTEBRAL AND SACRAL BODY METASTASIS

Metastatic malignant disease of the spine generally involves the vertebrae/sacrum, with pain being the most frequent complaint.

Treatment of Vertebral and Sacral Body Metastasis

While radiotherapy and chemotherapy are frequently effective in reducing tumor burden and associated symptoms, pain relief may be delayed days to weeks, depending on tumor response. Further, these approaches rely on bone remodeling to regain strength in the vertebrae/sacrum, which may necessitate supportive bracing to minimize the risk of vertebral/sacral collapse during healing. Improvements in pain and function are the primary outcomes of interest for treatment of bone malignancy with percutaneous vertebroplasty or sacroplasty.

SURGICAL TREATMENT OPTIONS

Percutaneous Vertebroplasty

Vertebroplasty is a surgical procedure that involves the injection of synthetic cement (e.g., polymethylmethacrylate, bis-glycidol dimethacrylate [Cortoss]³) into a fractured vertebra. It has been suggested that vertebroplasty may provide an analgesic effect through mechanical stabilization of a fractured or otherwise weakened vertebral body. However, other mechanisms of effect have been postulated, including thermal damage to intraosseous nerve fibers.

Percutaneous Sacroplasty

Sacroplasty evolved from the treatment of insufficiency fractures in the thoracic and lumbar vertebrae with vertebroplasty. The procedure, essentially identical to vertebroplasty, entails guided injection of polymethylmethacrylate through a needle inserted into the fracture zone. Although first described in 2000 as a treatment for symptomatic sacral metastatic lesions,^{4,5} it is most often described as a minimally invasive alternative to conservative management⁶⁻⁸ for sacral insufficiency fractures.

Pain and function are subjective outcomes and, thus, may be susceptible to placebo effects. Furthermore, the natural history of pain and disability associated with these conditions may vary. Therefore, controlled comparison studies would be valuable to demonstrate the clinical effectiveness of vertebroplasty and sacroplasty over and above any associated nonspecific or placebo effects and to demonstrate the effect of treatment compared with alternatives such as continued medical management.

In all clinical situations, adverse events related to complications from vertebroplasty and sacroplasty are the primary harms to be considered. Principal safety concerns relate to the incidence and consequences of leakage of the injected polymethyl methacrylate or another injectate.

REGULATORY STATUS

Vertebroplasty is a surgical procedure and, as such, is not subject to U.S. Food and Drug Administration (FDA) approval.

Polymethylmethacrylate bone cement was available as a drug product before enactment of the FDA's device regulation and was at first considered what the FDA terms a "transitional device." It was transitioned to a class III device requiring premarketing applications. Several orthopedic companies have received approval of their bone cement products since 1976. In 1999, polymethylmethacrylate was reclassified from class III to class II, which requires future 510(k) submissions to meet "special controls" instead of "general controls" to assure safety and effectiveness. Thus, use of polymethylmethacrylate in vertebroplasty represented an off-label use of an FDA-regulated product before 2005. In 2005, polymethylmethacrylate bone cements such as Spine-Fix[®] Biomimetic Bone Cement and Osteopal[®] V were cleared for marketing by the FDA through the 510(k) process for the fixation of pathologic fractures of the vertebral body using vertebroplasty procedures.

The use of polymethylmethacrylate in sacroplasty is an off-label use of an FDA-regulated product (bone cements such as Spine-Fix[®] Biomimetic Bone Cement [Teknimed] and Osteopal[®] V [Heraeus]) because the 510(k) approv-

al was for the fixation of pathologic fractures of the vertebral body using vertebroplasty procedures. Sacroplasty was not included. FDA product code: NDN.

In 2009, Cortoss® (Stryker) Bone Augmentation Material was cleared for marketing by the FDA through the 510(k) process. Cortoss® is a nonresorbable synthetic material that is a composite resin-based, bis-glycidyl dimethacrylate. The FDA classifies this product as a polymethylmethacrylate bone cement.

In 2010, the Parallax® Contour® Vertebral Augmentation Device (ArthroCare) was cleared for marketing by FDA through the 510(k) process. The device creates a void in cancellous bone that can then be filled with bone cement. FDA product code: HXG.

Services that are the subject of a clinical trial do not meet our Technology Assessment and Medically Necessary Services Protocol criteria and are considered investigational. *For explanation of experimental and investigational, please refer to the Technology Assessment and Medically Necessary Services Protocol.*

It is expected that only appropriate and medically necessary services will be rendered. We reserve the right to conduct prepayment and postpayment reviews to assess the medical appropriateness of the above-referenced procedures. **Some of this protocol may not pertain to the patients you provide care to, as it may relate to products that are not available in your geographic area.**

REFERENCES

We are not responsible for the continuing viability of web site addresses that may be listed in any references below.

1. Gotis-Graham I, McGuigan L, Diamond T, et al. Sacral insufficiency fractures in the elderly. *J Bone Joint Surg Br.* Nov 1994;76(6):882-886. PMID 7983111
2. Lin J, Lachmann E, Nagler W. Sacral insufficiency fractures: a report of two cases and a review of the literature. *J Womens Health Gend Based Med.* Sep 2001;10(7):699-705. PMID 11571100
3. Bae H, Hatten HP, Jr., Linovitz R, et al. A prospective randomized FDA-IDE trial comparing Cortoss with PMMA for vertebroplasty: a comparative effectiveness research study with 24-month follow-up. *Spine (Phila Pa 1976).* Apr 1 2012;37(7):544-550. PMID 21738093
4. Dehdashti AR, Martin JB, Jean B, et al. PMMA cementoplasty in symptomatic metastatic lesions of the S1 vertebral body. *Cardiovasc Intervent Radiol.* May-Jun 2000;23(3):235-237. PMID 10821903
5. Marcy PY, Palussiere J, Descamps B, et al. Percutaneous cementoplasty for pelvic bone metastasis. *Support Care Cancer.* Nov 2000;8(6):500-503. PMID 11094996
6. Aretxabala I, Fraiz E, Perez-Ruiz F, et al. Sacral insufficiency fractures. High association with pubic rami fractures. *Clin Rheumatol.* Oct 2000;19(5):399-401. PMID 11055834
7. Leroux JL, Denat B, Thomas E, et al. Sacral insufficiency fractures presenting as acute low-back pain. Biomechanical aspects. *Spine (Phila Pa 1976).* Dec 1993;18(16):2502-2506. PMID 8303454
8. Newhouse KE, el-Khoury GY, Buckwalter JA. Occult sacral fractures in osteopenic patients. *J Bone Joint Surg Am.* Dec 1992;74(10):1472-1477. PMID 1364816
9. Stratford PW, Binkley J, Solomon P, et al. Defining the minimum level of detectable change for the Roland-Morris questionnaire. *Phys Ther.* 1996 Apr;76(4):359-65. PMID: 8606899
10. Katz J, Melzack R. Measurement of pain. *Surg Clin North Am.* 1999 Apr;79(2):231-52. PMID: 10352653
11. Blue Cross and Blue Shield Technology Evaluation Center (TEC). Percutaneous Vertebroplasty. TEC Assessments. 2000;Volume 15:Tab 21.

12. Blue Cross and Blue Shield Technology Evaluation Center (TEC). Percutaneous vertebroplasty for vertebral fractures caused by osteoporosis, malignancy, or hemangioma. TEC Assessments. 2004;Volume 19:Tab 13.
13. Blue Cross and Blue Shield Technology Evaluation Center (TEC). Percutaneous vertebroplasty for vertebral fractures caused by osteoporosis or malignancy. TEC Assessments. 2005;Volume 20:Tab 6.
14. Blue Cross and Blue Shield Technology Evaluation Center (TEC). Percutaneous vertebroplasty or kyphoplasty for vertebral fractures caused by osteoporosis or malignancy. TEC Assessments. 2008;Volume 23:Tab 5.
15. Blue Cross and Blue Shield Technology Evaluation Center (TEC). Percutaneous vertebroplasty or kyphoplasty for vertebral fractures caused by osteoporosis. TEC Assessments. 2009;Volume 24:Tab 7.
16. Blue Cross and Blue Shield Technology Evaluation Center (TEC). Percutaneous vertebroplasty for vertebral fractures caused by osteoporosis. TEC Assessments. 2010;Volume 25:Tab 9.
17. Buchbinder R, Johnston RV, Rischin KJ, et al. Percutaneous vertebroplasty for osteoporotic vertebral compression fracture. *Cochrane Database Syst Rev*. 2018 Apr 4;4:CD006349. PMID: 29618171
18. Staples MP, Kallmes DF, Comstock BA, et al. Effectiveness of vertebroplasty using individual patient data from two randomised placebo controlled trials: meta-analysis. *BMJ*. Jul 12 2011;343:d3952. PMID 21750078
19. Xie L, Zhao ZG, Zhang SJ, et al. Percutaneous vertebroplasty versus conservative treatment for osteoporotic vertebral compression fractures: An updated meta-analysis of prospective randomized controlled trials. *Int J Surg*. Nov 2017;47:25-32. PMID 28939236
20. Buchbinder R, Osborne RH, Ebeling PR, et al. A randomized trial of vertebroplasty for painful osteoporotic vertebral fractures. *N Engl J Med*. Aug 6 2009;361(6):557-568. PMID 19657121
21. Kallmes DF, Comstock BA, Heagerty PJ, et al. A randomized trial of vertebroplasty for osteoporotic spinal fractures. *N Engl J Med*. Aug 6 2009;361(6):569-579. PMID 19657122
22. Kroon F, Staples M, Ebeling PR, et al. Two-year results of a randomized placebo-controlled trial of vertebroplasty for acute osteoporotic vertebral fractures. *J Bone Miner Res*. Jun 2014;29(6):1346-1355. PMID 24967454
23. Ostelo RW, Deyo RA, Stratford P, et al. Interpreting change scores for pain and functional status in low back pain: towards international consensus regarding minimal important change. *Spine (Phila Pa 1976)*. Jan 1 2008;33(1):90-94. PMID 18165753
24. Comstock BA, Sitlani CM, Jarvik JG, et al. Investigational vertebroplasty safety and efficacy trial (INVEST): patient-reported outcomes through 1 year. *Radiology*. Oct 2013;269(1):224-231. PMID 23696683
25. Firanescu CE, de Vries J, Lodder P, et al. Vertebroplasty versus sham procedure for painful acute osteoporotic vertebral compression fractures (VERTOS IV): randomised sham controlled clinical trial. *BMJ*. 2018 May 9;361:k1551. PMID: 29743284
26. Chen D, An ZQ, Song S, et al. Percutaneous vertebroplasty compared with conservative treatment in patients with chronic painful osteoporotic spinal fractures. *J Clin Neurosci*. Mar 2014;21(3):473-477. PMID 24315046
27. Farrokhi MR, Alibai E, Maghami Z. Randomized controlled trial of percutaneous vertebroplasty versus optimal medical management for the relief of pain and disability in acute osteoporotic vertebral compression fractures. *J Neurosurg Spine*. May 2011;14(5):561-569. PMID 21375382
28. Edidin AA, Ong KL, Lau E, et al. Mortality risk for operated and nonoperated vertebral fracture patients in the Medicare population. *J Bone Miner Res*. Jul 2011;26(7):1617-1626. PMID 21308780
29. Edidin AA, Ong KL, Lau E, et al. Morbidity and mortality after vertebral fractures: comparison of vertebral augmentation and nonoperative management in the Medicare population. *Spine (Phila Pa 1976)*. Aug 01 2015;40(15):1228-1241. PMID 26020845
30. Lin JH, Chien LN, Tsai WL, et al. Early vertebroplasty associated with a lower risk of mortality and respiratory failure in aged patients with painful vertebral compression fractures: a population-based cohort study in Taiwan. *Spine J*. Sep 2017;17(9):1310-1318. PMID 28483705
31. Clark W, Bird P, Gonski P, et al. Safety and efficacy of vertebroplasty for acute painful osteoporotic fractures (VAPOUR): a multicentre, randomised, double-blind, placebo-controlled trial. *Lancet*. Oct 01 2016; 388(10052):1408-1416. PMID 27544377

32. Klazen CA, Lohle PN, de Vries J, et al. Vertebroplasty versus conservative treatment in acute osteoporotic vertebral compression fractures (Vertos II): an open-label randomised trial. *Lancet*. Sep 25 2010;376(9746):1085-1092. PMID 20701962
33. Yi X, Lu H, Tian F, et al. Recompression in new levels after percutaneous vertebroplasty and kyphoplasty compared with conservative treatment. *Arch Orthop Trauma Surg*. Jan 2014;134(1):21-30. PMID 24287674
34. Leali PT, Solla F, Maestretti G, et al. Safety and efficacy of vertebroplasty in the treatment of osteoporotic vertebral compression fractures: a prospective multicenter international randomized controlled study. *Clin Cases Miner Bone Metab*. Sep-Dec 2016;13(3):234-236. PMID 28228788
35. Yang EZ, Xu JG, Huang GZ, et al. Percutaneous Vertebroplasty versus conservative treatment in aged patients with acute osteoporotic vertebral compression fractures: a prospective randomized controlled clinical study. *Spine (Phila Pa 1976)*. Apr 2016;41(8):653-660. PMID 26630417
36. Lourie H. Spontaneous osteoporotic fracture of the sacrum. An unrecognized syndrome of the elderly. *JAMA*. Aug 13 1982;248(6):715-717. PMID 7097924
37. Frey ME, Depalma MJ, Cifu DX, et al. Percutaneous sacroplasty for osteoporotic sacral insufficiency fractures: a prospective, multicenter, observational pilot study. *Spine J*. Mar-Apr 2008;8(2):367-373. PMID 17981097
38. Kortman K, Ortiz O, Miller T, et al. Multicenter study to assess the efficacy and safety of sacroplasty in patients with osteoporotic sacral insufficiency fractures or pathologic sacral lesions. *J Neurointerv Surg*. Sep 1 2013;5(5):461-466. PMID 22684691
39. Frey ME, Warner C, Thomas SM, et al. Sacroplasty: a ten-year analysis of prospective patients treated with percutaneous sacroplasty: literature review and technical considerations. *Pain Physician*. Nov 2017;20(7):E1063-E1072. PMID 29149151
40. Dougherty RW, McDonald JS, Cho YW, et al. Percutaneous sacroplasty using CT guidance for pain palliation in sacral insufficiency fractures. *J Neurointerv Surg*. Jan 2014;6(1):57-60. PMID 23345629
41. Zaman FM, Frey M, Slipman CW. Sacral stress fractures. *Curr Sports Med Rep*. Feb 2006;5(1):37-43. PMID 16483515
42. Denis F, Davis S, Comfort T. Sacral fractures: an important problem. Retrospective analysis of 236 cases. *Clin Orthop Relat Res*. Feb 1988;227:67-81. PMID 3338224
43. Shah LM, Jennings JW, Kirsch CFE et al. ACR Appropriateness Criteria(R) Management of Vertebral Compression Fractures. *J Am Coll Radiol*. 2018 Nov;15(11S). PMID 30392604
44. Barr JD, Jensen ME, Hirsch JA, et al. Position statement on percutaneous vertebral augmentation: a consensus statement developed by the Society of Interventional Radiology (SIR), American Association of Neurological Surgeons (AANS) and the Congress of Neurological Surgeons (CNS), American College of Radiology (ACR), American Society of Neuroradiology (ASNR), American Society of Spine Radiology (ASSR), Canadian Interventional Radiology Association (CIRA), and the Society of NeuroInterventional Surgery (SNIS). *J Vasc Interv Radiol*. 2014 Feb;25(2):171-81. PMID: 24325929.
45. Baerlocher MO, Saad WE, Dariushnia S, et al. Quality improvement guidelines for percutaneous vertebroplasty. *J Vasc Interv Radiol*. Feb 2014;25(2):165-170. PMID 24238815
46. McGuire R. AAOS Clinical Practice Guideline: the Treatment of Symptomatic Osteoporotic Spinal Compression Fractures. *J Am Acad Orthop Surg*. 2011 Mar;19(3). PMID 21368100
47. National Institute for Health and Care Excellence (NICE). Percutaneous vertebroplasty [IPG12]. 2003; <https://www.nice.org.uk/guidance/ipg12>. Accessed March 18, 2020.
48. National Institute for Health and Care Excellence (NICE). Percutaneous vertebroplasty and percutaneous balloon kyphoplasty for treating osteoporotic vertebral compression fractures [TA279]. 2013; <https://www.nice.org.uk/guidance/ta279>. Accessed March 18, 2020.
49. National Institute for Health and Care Excellence (NICE). Metastatic spinal cord compression in adults: risk assessment, diagnosis and management [CG75]. 2008; <https://www.nice.org.uk/guidance/cg75/chapter/1-guidance>. Accessed March 18, 2020.

50. National Government Services, Inc. (Primary Geographic Jurisdiction 06 & K - Illinois, Minnesota, Wisconsin, Connecticut, New York - Entire State, Maine, Massachusetts, New Hampshire, Rhode Island, Vermont) Local Coverage Determination (LCD): Percutaneous Vertebral Augmentation (PVA) for Osteoporotic Vertebral Compression Fracture (VCF) (L33569), Revision Effective Date for services performed on or after 12/01/2019.