

(701118)

Medical Benefit		Effective Date: 10/01/13	Next Review Date: 07/21
Preauthorization	No	Review Dates: 09/09, 09/10, 07/11, 07/12, 07/13, 07/14, 07/15, 07/16, 07/17, 07/18, 07/19, 07/20	

Preauthorization is not required.

The following protocol contains medical necessity criteria that apply for this service. The criteria are also applicable to services provided in the local Medicare Advantage operating area for those members, unless separate Medicare Advantage criteria are indicated. If the criteria are not met, reimbursement will be denied and the patient cannot be billed. Please note that payment for covered services is subject to eligibility and the limitations noted in the patient's contract at the time the services are rendered.

RELATED PROTOCOL

Hip Resurfacing

Populations	Interventions	Comparators	Outcomes
Individuals: <ul style="list-style-type: none"> Who are adults with asymptomatic femoroacetabular impingement 	Interventions of interest are: <ul style="list-style-type: none"> Femoroacetabular impingement surgery 	Comparators of interest are: <ul style="list-style-type: none"> Observation 	Relevant outcomes include: <ul style="list-style-type: none"> Symptoms Functional outcomes Health status measures Quality of life Change in disease status
Individuals: <ul style="list-style-type: none"> Who are adults with symptomatic femoroacetabular impingement 	Interventions of interest are: <ul style="list-style-type: none"> Femoroacetabular impingement surgery 	Comparators of interest are: <ul style="list-style-type: none"> Conservative management 	Relevant outcomes include: <ul style="list-style-type: none"> Symptoms Functional outcomes Health status measures Quality of life Change in disease status
Individuals: <ul style="list-style-type: none"> Who are children ≤15 years of age with symptomatic femoroacetabular impingement 	Interventions of interest are: <ul style="list-style-type: none"> Femoroacetabular impingement surgery 	Comparators of interest are: <ul style="list-style-type: none"> Conservative management 	Relevant outcomes include: <ul style="list-style-type: none"> Symptoms Functional outcomes Health status measures Quality of life Change in disease status
Individuals: <ul style="list-style-type: none"> Who are children ≤15 years of age with slipped capital femoral epiphysis-associated femoroacetabular impingement 	Interventions of interest are: <ul style="list-style-type: none"> Femoroacetabular impingement surgery 	Comparators of interest are: <ul style="list-style-type: none"> Surgical repair of slipped capital femoral epiphysis alone 	Relevant outcomes include: <ul style="list-style-type: none"> Symptoms Functional outcomes Health status measures Quality of life Change in disease status

Populations	Interventions	Comparators	Outcomes
Individuals: <ul style="list-style-type: none"> • With residual femoroacetabular impingement symptoms following primary surgery 	Interventions of interest are: <ul style="list-style-type: none"> • Revision arthroscopic surgery 	Comparators of interest are: <ul style="list-style-type: none"> • Conservative management 	Relevant outcomes include: <ul style="list-style-type: none"> • Symptoms • Functional outcomes • Health status measures • Quality of life • Change in disease status

DESCRIPTION

Femoroacetabular impingement results from localized compression within the joint as a result of an anatomic mismatch between the head of the femur and the acetabulum. Symptoms of impingement typically occur in young to middle-aged adults before the onset of osteoarthritis but may be present in younger patients with developmental hip disorders. The objective of surgical treatment of femoroacetabular impingement is to provide symptom relief and reduce further joint damage.

SUMMARY OF EVIDENCE

For individuals who are adults with asymptomatic femoroacetabular impingement who receive femoroacetabular impingement surgery, there is no direct evidence that the surgical treatment will prevent the development of osteoarthritis. Relevant outcomes are symptoms, functional outcomes, health status measures, quality of life, and change in disease status. Indirect evidence consists of observational studies. In retrospective studies of patients with osteoarthritis, the relevant outcomes were radiographic evidence of hip joint malformations. In prospective studies of patients with femoroacetabular impingement, the relevant outcome is progression to osteoarthritis. Several large observational studies (>1,000 patients), as well as smaller studies, have shown radiographic evidence of relationships between abnormal hip morphology and the development of osteoarthritis. There have been no studies in which femoroacetabular impingement surgery was performed on patients with femoroacetabular impingement morphology but no symptoms. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who are adults with symptomatic femoroacetabular impingement who receive femoroacetabular impingement surgery, the evidence includes systematic reviews of large and small observational studies and a small randomized controlled trial. Relevant outcomes are symptoms, functional outcomes, health status measures, quality of life, and change in disease status. Open hip dislocation surgery and arthroscopic surgery are the most common surgical techniques performed on patients with femoroacetabular impingement. Systematic reviews have evaluated open hip dislocation surgery and arthroscopic surgery, compared with no comparator, nonsurgical management, and other surgical techniques. Compared with nonsurgical management, all types of surgical techniques have resulted in significant improvements in functional outcomes, pain, and radiographic measurements. The reviews were limited when comparing surgical techniques with each other because patient characteristics and outcome measurements were heterogeneous among studies. The evidence is sufficient to determine the technology results in a meaningful improvement in the net health outcome.

For individuals who are children 15 years of age or younger with symptomatic femoroacetabular impingement who receive femoroacetabular impingement surgery, the evidence includes systematic reviews evaluating small observational studies and case series. Relevant outcomes are symptoms, functional outcomes, health status measures, quality of life, and change in disease status. While the studies reported reductions in pain and improvements in functional outcomes, the sample sizes were relatively small, with an average of 54 patients per study. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who are children 15 years of age or younger with slipped capital femoral epiphysis-associated femoroacetabular impingement who receive femoroacetabular impingement surgery, the evidence includes small observational studies (range, 19-51 patients). Relevant outcomes are symptoms, functional outcomes, health status measures, quality of life, and change in disease status. While most patients experienced symptom relief following femoroacetabular impingement surgery, the surgery is invasive and complications (e.g., non-unions) were reported. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have residual femoroacetabular impingement symptoms following a primary surgery who receive revision arthroscopic surgery, the evidence includes systematic reviews of observational studies (>400 patients). Relevant outcomes are symptoms, functional outcomes, health status measures, quality of life, and change in disease status. Though the studies were of low-quality, consistent improvements in functional outcomes, pain relief, and patient satisfaction were reported, in some cases beyond three years. The evidence is sufficient to determine the technology results in a meaningful improvement in the net health outcome.

POLICY

Open or arthroscopic treatment of femoroacetabular impingement (FAI) may be **medically necessary** when all of the following conditions have been met:

AGE

- Candidates should be skeletally mature with documented closure of growth plates (e.g., 15 years of age or older).

SYMPTOMS

1. Moderate-to-severe hip pain worsened by flexion activities (e.g., squatting or prolonged sitting) that significantly limits activities; AND
2. Unresponsive to conservative therapy for at least three months (including activity modifications, restriction of athletic pursuits and avoidance of symptomatic motion); AND
3. Positive impingement sign on clinical examination (pain elicited with 90° of flexion and internal rotation and adduction of the femur).

IMAGING

- Morphology indicative of cam or pincer FAI (e.g., pistol-grip deformity, femoral head-neck offset with an alpha angle greater than 50°, a positive wall sign, acetabular retroversion [overcoverage with crossover sign]), coxa profunda or protrusion, or damage of the acetabular rim; AND
- High probability of a causal association between the FAI morphology and damage (e.g., a pistol-grip deformity with a tear of the acetabular labrum and articular cartilage damage in the anterosuperior quadrant); AND
- No evidence of advanced osteoarthritis, defined as Tonnis grade 2 or 3, or joint space of less than two mm; AND
- No evidence of severe (Outerbridge grade IV) chondral damage.

Treatment of FAI is considered **investigational** in all other situations.

POLICY GUIDELINES

If FAI morphology is identified, patients should be advised not to play aggressive sports. It should be noted that current imaging techniques limit the early identification of cartilage defects, whereas delay in the surgical correction of bony abnormalities may lead to disease progression to the point at which joint preservation is no longer appropriate. In addition to magnetic resonance arthrography (MRA) some clinicians may also use local anesthetic injection into the joint to assist in confirming FAI pathology.

Treatment of FAI should be restricted to centers experienced in treating this condition and staffed by surgeons adequately trained in techniques addressing FAI. Because of the differing benefits and risks of open and arthroscopic approaches, patients should make an informed choice between the procedures.

Some patients may require a revision procedure if symptoms recur or persist. Published studies have indicated that all sources of impingement might not have been identified before surgery, and those that had might not have been adequately treated. The risk of needing an additional surgical procedure can be reduced by intra-operative assessment of impingement after bone debridement and reshaping.

BACKGROUND

FEMOROACETABULAR IMPINGEMENT

Femoroacetabular impingement arises from an anatomic mismatch between the head of the femur and the acetabulum, causing compression of the labrum or articular cartilage during flexion. The mismatch can arise from subtle morphologic alterations in the anatomy or orientation of the ball-and-socket components (e.g., a bony prominence at the head-neck junction or acetabular overcoverage), with articular cartilage damage initially occurring from abutment of the femoral neck against the acetabular rim, typically at the anterosuperior aspect of the acetabulum. Although hip joints can possess the morphologic features of femoroacetabular impingement without symptoms, femoroacetabular impingement may become pathologic with repetitive movement and/or increased force on the hip joint. High-demand activities may also result in pathologic impingement in hips with normal morphology.

Two types of impingement, cam, and pincer, may occur alone or, more frequently, together. Cam impingement is associated with an asymmetric or nonspherical contour of the head or neck of the femur jamming against the acetabulum, resulting in cartilage damage and delamination (detachment from the subchondral bone). Deformity of the head/neck junction that looks like a pistol-grip on radiographs is associated with damage to the anterosuperior area of the acetabulum. Symptomatic cam impingement is found most frequently in young male athletes. Pincer impingement is associated with overcoverage of the acetabulum and pinching of the labrum, with pain more typically beginning in women of middle age. In cases of isolated pincer impingement, the damage may be limited to a narrow strip of the acetabular cartilage.

Epidemiologic and radiographic studies have found correlations between hip osteoarthritis and femoroacetabular impingement lesions, supporting the theory that prolonged contact between the anatomically mismatched acetabulum and femur may lead not only to cam and pincer lesions but also to further cartilage damage and subsequent joint deterioration. It is believed that osteoplasty of the impinging bone is needed to protect the cartilage from further damage and to preserve the natural joint. Therefore, if femoroacetabular impingement morphology is shown to be an etiology of osteoarthritis, a strategy to reduce the occurrence of idiopathic hip osteoarthritis could be early recognition and treatment of femoroacetabular impingement before cartilage damage and joint deterioration occurs.

An association between femoroacetabular impingement and athletic pubalgia, sometimes called sports hernia, has been proposed. Athletic pubalgia is an umbrella term for a large variety of musculoskeletal injuries involving attachments and/or soft tissue support structures of the pubis.

Treatment

A technique for hip dislocation with open osteo chondroplasty that preserved the femoral blood supply was reported by Ganz. Visualization of the entire joint with this procedure led to the identification and acceptance of femoroacetabular impingement as an etiology of cartilage damage and the possibility of correcting the abnormal femoroacetabular morphology. Open osteo chondroplasty of bony abnormalities and treatment of the symptomatic cartilage defect is considered the criterion standard for complex bony abnormalities. However, open osteo chondroplasty is invasive, requiring transection of the greater trochanter (separation of the femoral head from the femoral shaft) and dislocation of the hip joint to provide full access to the femoral head and acetabulum. In addition to the general adverse events of open surgical procedures, open osteo chondroplasty with dislocation has been associated with nonunion and neurologic and soft tissue lesions.

Less invasive hip arthroscopy and an arthroscopy-assisted mini-approach were developed by 2004. Arthroscopy requires specially designed instruments and is considered technically more difficult due to reduced visibility and limited access to the joint space. Advanced imaging techniques, including computed tomography and fluoroscopy, have been used to improve visualization of the three-dimensional head/neck morphology during arthroscopy.

Femoroacetabular impingement can also be a source of hip pain and decreased hip internal rotation in the pediatric population. When nonoperative management of femoroacetabular impingement in children and adolescents is ineffective, surgical procedures may be indicated. Surgical techniques include arthroscopy, open hip dislocation, limited open with arthroscopy, and osteotomy.

SLIPPED CAPITAL FEMORAL EPIPHYSIS

Patients with slipped capital femoral epiphysis have a displaced femoral head in relation to the femoral neck within the confines of the acetabulum, which can result in hip pain, thigh pain, knee pain, and the onset of a limp. Slipped capital femoral epiphysis occurs most frequently in children between the ages of 10 to 16. Upon reaching skeletal maturity patients diagnosed with slipped capital femoral epiphysis, 32% were found to have clinical signs of impingement. It is not uncommon for patients with slipped capital femoral epiphysis to develop premature osteoarthritis and require total hip arthroplasty within 20 years.

Treatment

The standard treatment for slipped capital femoral epiphysis is stabilization across the physis by in situ pinning. Alternative treatments proposed for pediatric patients with slipped capital femoral epiphysis related femoroacetabular impingement include osteoplasty without dislocation, or with the open dislocation technique described by Ganz. The Ganz technique (capital realignment with open dislocation) is technically demanding, with a steep learning curve and a high-risk of complications, including avascular necrosis. Therefore, early treatment to decrease impingement must be weighed against the increased risk of adverse events.

REGULATORY STATUS

Surgery for treatment of femoroacetabular impingement is a procedure and, as such, is not subject to regulation by the U.S. Food and Drug Administration.

Services that are the subject of a clinical trial do not meet our Technology Assessment and Medically Necessary Services Protocol criteria and are considered investigational. *For explanation of experimental and investigational, please refer to the Technology Assessment and Medically Necessary Services Protocol.*

It is expected that only appropriate and medically necessary services will be rendered. We reserve the right to conduct prepayment and postpayment reviews to assess the medical appropriateness of the above-referenced procedures. **Some of this protocol may not pertain to the patients you provide care to, as it may relate to products that are not available in your geographic area.**

REFERENCES

We are not responsible for the continuing viability of web site addresses that may be listed in any references below.

1. Egger AC, Frangiamore S, Rosneck J. Femoroacetabular Impingement: A Review. *Sports Med Arthrosc Rev.* 2016 Dec;24(4). PMID 27811519
2. Frank JM, Harris JD, Erickson BJ et al. Prevalence of Femoroacetabular Impingement Imaging Findings in Asymptomatic Volunteers: A Systematic Review. *Arthroscopy.* 2015 Jun;31(6). PMID 25636988
3. Oner A, Koksak A, Sofu H, et al. The prevalence of femoroacetabular impingement as an aetiological factor for end-stage degenerative osteoarthritis of the hip joint: analysis of 1,000 cases. *Hip Int.* Mar-Apr 2016;26(2): 164-168. PMID 26916653
4. Thomas GE, Palmer AJ, Batra RN, et al. Subclinical deformities of the hip are significant predictors of radiographic osteoarthritis and joint replacement in women. A 20 year longitudinal cohort study. *Osteoarthritis Cartilage.* Oct 2014;22(10):1504-1510. PMID 25047637
5. Reichenbach S, Leunig M, Werlen S, et al. Association between cam-type deformities and magnetic resonance imaging-detected structural hip damage: a cross-sectional study in young men. *Arthritis Rheum.* Dec 2011;63(12):4023-4030. PMID 21904996
6. Gosvig KK, Jacobsen S, Sonne-Holm S, et al. Prevalence of malformations of the hip joint and their relationship to sex, groin pain, and risk of osteoarthritis: a population-based survey. *J Bone Joint Surg Am.* May 2010;92(5):1162-1169. PMID 20439662
7. Takeyama A, Naito M, Shiramizu K, et al. Prevalence of femoroacetabular impingement in Asian patients with osteoarthritis of the hip. *Int Orthop.* Oct 2009;33(5):1229-1232. PMID 19277653
8. Bardakos NV, Villar RN. Predictors of progression of osteoarthritis in femoroacetabular impingement: a radiological study with a minimum of ten years follow-up. *J Bone Joint Surg Br.* Feb 2009;91(2):162-169. PMID 19190047
9. Kim KC, Hwang DS, Lee CH, et al. Influence of femoroacetabular impingement on results of hip arthroscopy in patients with early osteoarthritis. *Clin Orthop Relat Res.* Mar 2007;456:128-132. PMID 17106273
10. Beck M, Kalhor M, Leunig M, et al. Hip morphology influences the pattern of damage to the acetabular cartilage: femoroacetabular impingement as a cause of early osteoarthritis of the hip. *J Bone Joint Surg Br.* Jul 2005;87(7):1012-1018. PMID 15972923
11. Tanzer M, Noiseux N. Osseous abnormalities and early osteoarthritis: the role of hip impingement. *Clin Orthop Relat Res.* Dec 2004(429):170-177. PMID 15577483
12. Reiman MP, Peters S, Sylvain J et al. Femoroacetabular impingement surgery allows 74% of athletes to return to the same competitive level of sports participation but their level of performance remains unreported: a systematic review with meta-analysis. *Br J Sports Med.* 2018 Aug;52(15). PMID 29581142
13. Wall PD, Brown JS, Parsons N, et al. Surgery for treating hip impingement (femoroacetabular impingement). *Cochrane Database Syst Rev.* Sep 8 2014;9(9):CD010796. PMID 25198064
14. Harris JD, Erickson BJ, Bush-Joseph CA, et al. Treatment of femoroacetabular impingement: a systematic review. *Curr Rev Musculoskelet Med.* Sep 2013;6(3):207-218. PMID 23743861
15. Bedi A, Chen N, Robertson W, et al. The management of labral tears and femoroacetabular impingement of the hip in the young, active patient. *Arthroscopy.* Oct 2008;24(10):1135-1145. PMID 19028166
16. Espinosa N, Rothenfluh DA, Beck M, et al. Treatment of femoro-acetabular impingement: preliminary results of labral refixation. *J Bone Joint Surg Am.* May 2006;88(5):925-935. PMID 16651565

17. Peters CL, Erickson JA. Treatment of femoro-acetabular impingement with surgical dislocation and debridement in young adults. *J Bone Joint Surg Am.* Aug 2006;88(8):1735-1741. PMID 16882895
18. Beck M, Leunig M, Parvizi J, et al. Anterior femoroacetabular impingement: part II. Midterm results of surgical treatment. *Clin Orthop Relat Res.* Jan 2004(418):67-73. PMID 15043095
19. Dwyer T, Whelan D, Shah PS et al. Operative Versus Nonoperative Treatment of Femoroacetabular Impingement Syndrome: A Meta-analysis of Short-Term Outcomes. *Arthroscopy.* 2020 Jan;36(1). PMID 31864588
20. Minkara AA, Westermann RW, Rosneck J, et al. Systematic review and meta-analysis of outcomes after hip arthroscopy in femoroacetabular impingement. *Am J Sports Med.* Jan 1 2018;363546517749475. PMID 29373805
21. Kierkegaard S, Langeskov-Christensen M, Lund B, et al. Pain, activities of daily living and sport function at different time points after hip arthroscopy in patients with femoroacetabular impingement: a systematic review with meta-analysis. *Br J Sports Med.* Apr 2017;51(7):572-579. PMID 27845683
22. Krych AJ, Thompson M, Knutson Z, et al. Arthroscopic labral repair versus selective labral debridement in female patients with femoroacetabular impingement: a prospective randomized study. *Arthroscopy.* Jan 2013;29(1):46-53. PMID 23276413
23. Palmer AJR, Ayyar Gupta V, Fernquest S, et al. *BMJ.* 2019 Feb 7;364:l185. PMID: 30733197
24. Griffin DR, Dickenson EJ, Wall PDH, et al. Hip arthroscopy versus best conservative care for the treatment of femoroacetabular impingement syndrome (UK FASHIoN): a multicenter randomised controlled trial. *Lancet.* 2018 Jun 2;391(10136):2225-2235. PMID: 29893223
25. Zhang D, Chen L, Wang G. Hip arthroscopy versus open surgical dislocation for femoroacetabular impingement: A systematic review and meta-analysis. *Medicine (Baltimore).* Oct 2016;95(41):e5122. PMID 27741133
26. Nwachukwu BU, Rebolledo BJ, McCormick F, et al. Arthroscopic versus open treatment of femoroacetabular impingement: a systematic review of medium- to long-term outcomes. *Am J Sports Med.* Apr 2016;44(4):-1062-1068. PMID 26059179
27. Matsuda DK, Carlisle JC, Arthurs SC, et al. Comparative systematic review of the open dislocation, mini-open, and arthroscopic surgeries for femoroacetabular impingement. *Arthroscopy.* Feb 2011;27(2):252-269. PMID 21266276
28. Botser IB, Smith TW, Jr., Nasser R, et al. Open surgical dislocation versus arthroscopy for femoroacetabular impingement: a comparison of clinical outcomes. *Arthroscopy.* Feb 2011;27(2):270-278. PMID 21266277
29. Papalia R, Del Buono A, Franceschi F, et al. Femoroacetabular impingement syndrome management: Arthroscopy or open surgery? *Int Orthop.* May 2012;36(5):903-914. PMID 22190060
30. Zingg PO, Ulbrich EJ, Buehler TC, et al. Surgical hip dislocation versus hip arthroscopy for femoroacetabular impingement: clinical and morphological short-term results. *Arch Orthop Trauma Surg.* Jan 2013;133(1):69-79. PMID 23064993
31. Domb BG, Stake CE, Botser IB, et al. Surgical dislocation of the hip versus arthroscopic treatment of femoroacetabular impingement: a prospective matched-pair study with average 2-year follow-up. *Arthroscopy.* Sep 2013;29(9):1506-1513. PMID 23992988
32. Wu CT, Mahameed M, Lin PC et al. Treatment of cam-type femoroacetabular impingement using anterolateral mini-open and arthroscopic osteochondroplasty. *J Orthop Surg Res.* 2019 Jul;14(1). PMID 31315654
33. Chiron P, Espie A, Reina N, et al. Surgery for femoroacetabular impingement using a minimally invasive anterolateral approach: analysis of 118 cases at 2.2-year follow-up. *Orthop Traumatol Surg Res.* Feb 2012; 98(1):30-38. PMID 22257764
34. Laude F, Soriali E, Nogier A. Femoroacetabular impingement treatment using arthroscopy and anterior approach. *Clin Orthop Relat Res.* Mar 2009; 467(3):747-52. PMID 19089524
35. Oduwole KO, de Sa D, Kay J, et al. Surgical treatment of femoroacetabular impingement following slipped capital femoral epiphysis: A systematic review. *Bone Joint Res.* Aug 2017;6(8):472-480. PMID 28790036

36. de Sa D, Cargnelli S, Catapano M, et al. Femoroacetabular impingement in skeletally immature patients: a systematic review examining indications, outcomes, and complications of open and arthroscopic treatment. *Arthroscopy*. Feb 2015;31(2):373-384. PMID 25262968
37. Guindani N, Eberhardt O, Wirth T, et al. Surgical dislocation for pediatric and adolescent hip deformity: clinical and radiographical results at 3 years follow-up. *Arch Orthop Trauma Surg*. Apr 2017;137(4):471-479. PMID 28197752
38. Nwachukwu BU, Chang B, Kahlenberg CA, et al. Arthroscopic treatment of femoroacetabular impingement in adolescents provides clinically significant outcome improvement. *Arthroscopy*. Oct 2017;33(10):1812-1818. PMID 28623078
39. Tran P, Pritchard M, O'Donnell J. Outcome of arthroscopic treatment for cam type femoroacetabular impingement in adolescents. *ANZ J Surg*. Sep 3 2013;83(5):382-386. PMID 22943465
40. Sink EL, Zaltz I, Heare T, et al. Acetabular cartilage and labral damage observed during surgical hip dislocation for stable slipped capital femoral epiphysis. *J Pediatr Orthop*. Jan-Feb 2010;30(1):26-30. PMID 20032738
41. Ziebarth K, Zilkens C, Spencer S, et al. Capital realignment for moderate and severe SCFE using a modified Dunn procedure. *Clin Orthop Relat Res*. Mar 2009;467(3):704-716. PMID 19142692
42. Spencer S, Millis MB, Kim YJ. Early results of treatment of hip impingement syndrome in slipped capital femoral epiphysis and pistol grip deformity of the femoral head-neck junction using the surgical dislocation technique. *J Pediatr Orthop*. May-Jun 2006;26(3):281-285. PMID 16670535
43. O'Connor M, Steinl GK, Padaki AS et al. Outcomes of Revision Hip Arthroscopic Surgery: A Systematic Review and Meta-analysis. *Am J Sports Med*. 2019 Sep;363546519869671. PMID 31503501
44. Sardana V, Philippon MJ, de Sa D, et al. Revision hip arthroscopy indications and outcomes: a systematic review. *Arthroscopy*. Oct 2015;31(10):2047-2055. PMID 26033461
45. Cvetanovich GL, Harris JD, Erickson BJ, et al. Revision hip arthroscopy: a systematic review of diagnoses, operative findings, and outcomes. *Arthroscopy*. Jul 2015;31(7):1382-1390. PMID 25703289
46. Gwathmey FW, Jones KS, Thomas Byrd JW. Revision hip arthroscopy: findings and outcomes. *J Hip Preserv Surg*. Dec 2017;4(4):318-323. PMID 29250340
47. Newman JT, Briggs KK, McNamara SC, et al. Outcomes after revision hip arthroscopic surgery in adolescent patients compared with a matched cohort undergoing primary arthroscopic surgery. *Am J Sports Med*. Dec 2016;44(12):3063-3069. PMID 27514736
48. National Institute for Health and Clinical Excellence (NICE). Arthroscopic femoroacetabular surgery for hip impingement syndrome [IPG408]. 2011; <https://www.nice.org.uk/guidance/IPG408>. Accessed March 10, 2020.
49. National Institute for Health and Care Excellence (NICE). Open femoro-acetabular surgery for hip impingement syndrome [IPG403]. 2011; <https://www.nice.org.uk/guidance/IPG403>. Accessed March 10, 2020.