

Backward Map No Dx

I10_code_desc	Short Descriptor	I9 Map	Description
R40.2130	Coma scale, eyes open, to sound, unspecified time	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. And the vast majority of codes on this map are for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2131	Coma scale, eyes open, to sound, in the field	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2132	Coma scale, eyes open, to sound, EMR	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2133	Coma scale, eyes open, to sound, at hospital admission	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2134	Coma scale, eyes open, to sound, 24+hrs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2140	Coma scale, eyes open, spontaneous, unspecified time	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

R40.2141	Coma scale, eyes open, spontaneous, in the field	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2142	Coma scale, eyes open, spontaneous, EMR	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2143	Coma scale, eyes open, spontaneous, at hospital admission	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2144	Coma scale, eyes open, spontaneous, 24+hurs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2230	Coma scale, best verb, inappropriate words, unsp time	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2231	Coma scale, best verb, inappropriate words, in the field	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2232	Coma scale, best verbal response, inappropriate words, EMR	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

R40.2233	Coma scale, best verbal response, inappropriate words, admit	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2234	Coma scale, best verb, inappropriate words, 24+hrs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2240	Coma scale, best verb, confused conversation, unsp time	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2241	Coma scale, best verb, confused conversation, in the field	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2242	Coma scale, best verbal response, confused conversation, EMR	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2243	Coma scale, best verb, confused conversation, admit	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2244	Coma scale, best verb, confused conversation, 24+hrs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

R40.2250	Coma scale, best verbal response, oriented, unspecified time	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2251	Coma scale, best verbal response, oriented, in the field	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2252	Coma scale, best verbal response, oriented, EMR	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2253	Coma scale, best verbal response, oriented, admit	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2254	Coma scale, best verbal response, oriented, 24+hrs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2330	Coma scale, best motor response, abnormal, unspecified time	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2331	Coma scale, best motor response, abnormal, in the field	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

R40.2332	Coma scale, best motor response, abnormal, EMR	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2333	Coma scale, best motor response, abnormal, admit	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2334	Coma scale, best motor response, abnormal, 24+hrs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2350	Coma scale, best motor response, localizes pain, unsp time	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2351	Coma scale, best motor, localizes pain, in the field	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2352	Coma scale, best motor response, localizes pain, EMR	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2353	Coma scale, best motor response, localizes pain, admit	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

R40.2354	Coma scale, best motor response, localizes pain, 24+hrs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2360	Coma scale, best motor response, obeys commands, unsp time	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2361	Coma scale, best motor, obeys commands, in the field	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2362	Coma scale, best motor response, obeys commands, EMR	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2363	Coma scale, best motor response, obeys commands, admit	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.2364	Coma scale, best motor response, obeys commands, 24+hrs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.241	Glasgow coma scale score 13-15	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

R40.242	Glasgow coma scale score 9-12	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.243	Glasgow coma scale score 3-8	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
R40.244	Oth coma,w/o Glasgow coma scale score,or w/part score report	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. The coma scale codes are configured completely differently in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T36.0X6A	Underdosing of penicillins, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T36.0X6D	Underdosing of penicillins, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T36.0X6S	Underdosing of penicillins, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T36.1X6A	Underdosing of cephalospor/oth beta-lactm antibiotics, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T36.1X6D	Underdosing of cephalospor/oth beta-lactm antibiotics, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T36.1X6S	Underdosing of cephalospor/oth beta-lactm antibiot, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T36.2X6A	Underdosing of chloramphenicol group, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T36.2X6D	Underdosing of chloramphenicol group, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T36.2X6S	Underdosing of chloramphenicol group, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T36.3X6A	Underdosing of macrolides, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T36.3X6D	Underdosing of macrolides, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T36.3X6S	Underdosing of macrolides, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T36.4X6A	Underdosing of tetracyclines, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T36.4X6D	Underdosing of tetracyclines, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T36.4X6S	Underdosing of tetracyclines, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T36.5X6A	Underdosing of aminoglycosides, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T36.5X6D	Underdosing of aminoglycosides, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T36.5X6S	Underdosing of aminoglycosides, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T36.6X6A	Underdosing of rifampicins, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T36.6X6D	Underdosing of rifampicins, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T36.6X6S	Underdosing of rifampicins, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T36.7X6A	Underdosing of antifungal antibiotics, sys used, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T36.7X6D	Underdosing of antifungal antibiotics, sys used, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T36.7X6S	Underdosing of antifungal antibiotics, sys used, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T36.8X6A	Underdosing of other systemic antibiotics, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T36.8X6D	Underdosing of other systemic antibiotics, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T36.8X6S	Underdosing of other systemic antibiotics, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T36.96XA	Underdosing of unspecified systemic antibiotic, init encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T36.96XD	Underdosing of unspecified systemic antibiotic, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T36.96XS	Underdosing of unspecified systemic antibiotic, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T37.0X6A	Underdosing of sulfonamides, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T37.0X6D	Underdosing of sulfonamides, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T37.0X6S	Underdosing of sulfonamides, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T37.1X6A	Underdosing of antimycobacterial drugs, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T37.1X6D	Underdosing of antimycobacterial drugs, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T37.1X6S	Underdosing of antimycobacterial drugs, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T37.2X6A	Underdosing of antimalari/drugs acting on bld protzoa, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T37.2X6D	Underdosing of antimalari/drugs acting on bld protzoa, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T37.2X6S	Undrdose of antimalari/drugs acting on bld protzoa, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T37.3X6A	Underdosing of other antiprotozoal drugs, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T37.3X6D	Underdosing of other antiprotozoal drugs, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T37.3X6S	Underdosing of other antiprotozoal drugs, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T37.4X6A	Underdosing of anthelmintics, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T37.4X6D	Underdosing of anthelmintics, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T37.4X6S	Underdosing of anthelmintics, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T37.5X6A	Underdosing of antiviral drugs, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T37.5X6D	Underdosing of antiviral drugs, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T37.5X6S	Underdosing of antiviral drugs, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T37.8X6A	Underdosing of systemic anti-infect/parasit, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T37.8X6D	Underdosing of systemic anti-infect/parasit, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T37.8X6S	Underdosing of systemic anti-infect/parasit, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T37.96XA	Underdosing of unsp systemic anti-infect/parasit, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T37.96XD	Underdosing of unsp systemic anti-infect/parasit, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T37.96XS	Underdosing of unsp systemic anti-infect/parasit, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.0X6A	Underdosing of glucocorticoids and synthetic analogues, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.0X6D	Underdosing of glucocorticoids and synthetic analogues, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.0X6S	Underdosing of glucocort/synth analog, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.1X6A	Underdosing of thyroid hormones and substitutes, init encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.1X6D	Underdosing of thyroid hormones and substitutes, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.1X6S	Underdosing of thyroid hormones and substitutes, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T38.2X6A	Underdosing of antithyroid drugs, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.2X6D	Underdosing of antithyroid drugs, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.2X6S	Underdosing of antithyroid drugs, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.3X6A	Underdosing of insulin and oral hypoglycemic drugs, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.3X6D	Underdosing of insulin and oral hypoglycemic drugs, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.3X6S	Underdosing of insulin and oral hypoglycemic drugs, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.4X6A	Underdosing of oral contraceptives, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T38.4X6D	Underdosing of oral contraceptives, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.4X6S	Underdosing of oral contraceptives, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.5X6A	Underdosing of other estrogens and progestogens, init encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.5X6D	Underdosing of other estrogens and progestogens, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.5X6S	Underdosing of other estrogens and progestogens, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.6X6A	Underdosing of antigonadtr/antiestr/antiandrg, NEC, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.6X6D	Underdosing of antigonadtr/antiestr/antiandrg, NEC, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T38.6X6S	Underdosing of antigonadtr/antiestr/antiandrg, NEC, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.7X6A	Underdosing of androgens and anabolic congeners, init encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.7X6D	Underdosing of androgens and anabolic congeners, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.7X6S	Underdosing of androgens and anabolic congeners, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.806A	Underdosing of unsp hormones and synthetic substitutes, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.806D	Underdosing of unsp hormones and synthetic substitutes, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.806S	Underdosing of unsp hormones and synthetic sub, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T38.816A	Underdosing of anterior pituitary hormones, init encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.816D	Underdosing of anterior pituitary hormones, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.816S	Underdosing of anterior pituitary hormones, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.896A	Underdosing of oth hormones and synthetic substitutes, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.896D	Underdosing of oth hormones and synthetic substitutes, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.896S	Underdosing of hormones and synthetic substitutes, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.906A	Underdosing of unspecified hormone antagonists, init encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T38.906D	Underdosing of unspecified hormone antagonists, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.906S	Underdosing of unspecified hormone antagonists, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.996A	Underdosing of other hormone antagonists, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.996D	Underdosing of other hormone antagonists, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T38.996S	Underdosing of other hormone antagonists, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T39.016A	Underdosing of aspirin, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T39.016D	Underdosing of aspirin, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T39.016S	Underdosing of aspirin, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T39.096A	Underdosing of salicylates, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T39.096D	Underdosing of salicylates, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T39.096S	Underdosing of salicylates, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T39.1X6A	Underdosing of 4-Aminophenol derivatives, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T39.1X6D	Underdosing of 4-Aminophenol derivatives, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T39.1X6S	Underdosing of 4-Aminophenol derivatives, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T39.2X6A	Underdosing of pyrazolone derivatives, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T39.2X6D	Underdosing of pyrazolone derivatives, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T39.2X6S	Underdosing of pyrazolone derivatives, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T39.316A	Underdosing of propionic acid derivatives, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T39.316D	Underdosing of propionic acid derivatives, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T39.316S	Underdosing of propionic acid derivatives, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T39.396A	Underdosing of nonsteroidal anti-inflammatory drugs, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T39.396D	Underdosing of nonsteroidal anti-inflammatory drugs, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T39.396S	Underdosing of nonsteroidal anti-inflammatory drugs, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T39.4X6A	Underdosing of antirheumatics, NEC, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T39.4X6D	Underdosing of antirheumatics, NEC, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T39.4X6S	Underdosing of antirheumatics, NEC, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T39.8X6A	Underdosing of nonopioid analges/antipyret, NEC, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T39.8X6D	Underdosing of nonopioid analges/antipyret, NEC, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T39.8X6S	Underdosing of nonopioid analges/antipyret, NEC, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T39.96XA	Underdosing of unsp nonopi analgs/antipyr/antirheu, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T39.96XD	Underdosing of unsp nonopi analgs/antipyr/antirheu, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T39.96XS	Underdosing of unsp nonopi analgs/antipyr/antirheu, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T40.0X6A	Underdosing of opium, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T40.0X6D	Underdosing of opium, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T40.0X6S	Underdosing of opium, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T40.2X6A	Underdosing of other opioids, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T40.2X6D	Underdosing of other opioids, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T40.2X6S	Underdosing of other opioids, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T40.3X6A	Underdosing of methadone, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T40.3X6D	Underdosing of methadone, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T40.3X6S	Underdosing of methadone, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T40.4X6A	Underdosing of other synthetic narcotics, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T40.4X6D	Underdosing of other synthetic narcotics, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T40.4X6S	Underdosing of other synthetic narcotics, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T40.5X6A	Underdosing of cocaine, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T40.5X6D	Underdosing of cocaine, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T40.5X6S	Underdosing of cocaine, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T40.606A	Underdosing of unspecified narcotics, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T40.606D	Underdosing of unspecified narcotics, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T40.606S	Underdosing of unspecified narcotics, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T40.696A	Underdosing of other narcotics, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T40.696D	Underdosing of other narcotics, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T40.696S	Underdosing of other narcotics, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T40.7X6A	Underdosing of cannabis (derivatives), initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T40.7X6D	Underdosing of cannabis (derivatives), subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T40.7X6S	Underdosing of cannabis (derivatives), sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T40.906A	Underdosing of unspecified psychodysleptics, init encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T40.906D	Underdosing of unspecified psychodysleptics, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T40.906S	Underdosing of unspecified psychodysleptics, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T40.996A	Underdosing of other psychodysleptics, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T40.996D	Underdosing of other psychodysleptics, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T40.996S	Underdosing of other psychodysleptics, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T41.0X6A	Underdosing of inhaled anesthetics, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T41.0X6D	Underdosing of inhaled anesthetics, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T41.0X6S	Underdosing of inhaled anesthetics, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T41.1X6A	Underdosing of intravenous anesthetics, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T41.1X6D	Underdosing of intravenous anesthetics, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T41.1X6S	Underdosing of intravenous anesthetics, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T41.206A	Underdosing of unspecified general anesthetics, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T41.206D	Underdosing of unspecified general anesthetics, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T41.206S	Underdosing of unspecified general anesthetics, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T41.296A	Underdosing of other general anesthetics, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T41.296D	Underdosing of other general anesthetics, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T41.296S	Underdosing of other general anesthetics, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T41.3X6A	Underdosing of local anesthetics, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T41.3X6D	Underdosing of local anesthetics, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T41.3X6S	Underdosing of local anesthetics, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T41.46XA	Underdosing of unspecified anesthetics, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T41.46XD	Underdosing of unspecified anesthetics, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T41.46XS	Underdosing of unspecified anesthetics, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T41.5X6A	Underdosing of therapeutic gases, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T41.5X6D	Underdosing of therapeutic gases, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T41.5X6S	Underdosing of therapeutic gases, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T42.0X6A	Underdosing of hydantoin derivatives, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T42.0X6D	Underdosing of hydantoin derivatives, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T42.0X6S	Underdosing of hydantoin derivatives, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T42.1X6A	Underdosing of iminostilbenes, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T42.1X6D	Underdosing of iminostilbenes, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T42.1X6S	Underdosing of iminostilbenes, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T42.2X6A	Underdosing of succinimides and oxazolidinediones, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T42.2X6D	Underdosing of succinimides and oxazolidinediones, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T42.2X6S	Underdosing of succinimides and oxazolidinediones, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T42.3X6A	Underdosing of barbiturates, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T42.3X6D	Underdosing of barbiturates, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T42.3X6S	Underdosing of barbiturates, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T42.4X6A	Underdosing of benzodiazepines, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T42.4X6D	Underdosing of benzodiazepines, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T42.4X6S	Underdosing of benzodiazepines, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T42.5X6A	Underdosing of mixed antiepileptics, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T42.5X6D	Underdosing of mixed antiepileptics, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T42.5X6S	Underdosing of mixed antiepileptics, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T42.6X6A	Underdosing of antiepileptic and sed-hypntc drugs, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T42.6X6D	Underdosing of antiepileptic and sed-hypntc drugs, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T42.6X6S	Underdosing of antiepileptic and sed-hypntc drugs, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T42.76XA	Underdosing of unsp antiepileptic and sed-hypntc drugs, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T42.76XD	Underdosing of unsp antiepileptic and sed-hypntc drugs, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T42.76XS	Underdosing of unsp antieplptc and sed-hypntc drugs, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T42.8X6A	Underdosing of antiparkns drug/centr muscle-tone depr, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T42.8X6D	Underdosing of antiparkns drug/centr muscle-tone depr, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T42.8X6S	Underdosing of antiparkns drug/centr musc-tone depr, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.016A	Underdosing of tricyclic antidepressants, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.016D	Underdosing of tricyclic antidepressants, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T43.016S	Underdosing of tricyclic antidepressants, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.026A	Underdosing of tetracyclic antidepressants, init encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.026D	Underdosing of tetracyclic antidepressants, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.026S	Underdosing of tetracyclic antidepressants, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.1X6A	Underdosing of MAO inhib antidepressants, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.1X6D	Underdosing of MAO inhib antidepressants, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.1X6S	Underdosing of MAO inhib antidepressants, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T43.206A	Underdosing of unspecified antidepressants, init encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.206D	Underdosing of unspecified antidepressants, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.206S	Underdosing of unspecified antidepressants, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.216A	Undrdose of selective seroton/norepineph reup inhibtr, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.216D	Undrdose of selective seroton/norepineph reup inhibtr, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.216S	Undrdose of slctv seroton/norepineph reup inhibtr, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.226A	Underdosing of selective serotonin reuptake inhibitors, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T43.226D	Underdosing of selective serotonin reuptake inhibitors, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.226S	Undrdose of selective serotonin reuptake inhibitors, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.296A	Underdosing of other antidepressants, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.296D	Underdosing of other antidepressants, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.296S	Underdosing of other antidepressants, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.3X6A	Underdosing of phenothiazine antipsychot/neurolept, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.3X6D	Underdosing of phenothiazine antipsychot/neurolept, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T43.3X6S	Underdosing of phenothiazine antipsychot/neurolept, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.4X6A	Underdosing of butyrophen/thiothixen neuroleptics, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.4X6D	Underdosing of butyrophen/thiothixen neuroleptics, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.4X6S	Underdosing of butyrophen/thiothixen neuroleptics, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.506A	Underdosing of unsp antipsychotics and neuroleptics, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.506D	Underdosing of unsp antipsychotics and neuroleptics, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.506S	Underdosing of unsp antipsychotics and neuroleptics, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T43.596A	Underdosing of oth antipsychotics and neuroleptics, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.596D	Underdosing of oth antipsychotics and neuroleptics, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.596S	Underdosing of oth antipsychotics and neuroleptics, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.606A	Underdosing of unspecified psychostimulants, init encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.606D	Underdosing of unspecified psychostimulants, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.606S	Underdosing of unspecified psychostimulants, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.616A	Underdosing of caffeine, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T43.616D	Underdosing of caffeine, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.616S	Underdosing of caffeine, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.626A	Underdosing of amphetamines, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.626D	Underdosing of amphetamines, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.626S	Underdosing of amphetamines, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.636A	Underdosing of methylphenidate, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.636D	Underdosing of methylphenidate, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T43.636S	Underdosing of methylphenidate, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.696A	Underdosing of other psychostimulants, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.696D	Underdosing of other psychostimulants, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.696S	Underdosing of other psychostimulants, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.8X6A	Underdosing of other psychotropic drugs, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.8X6D	Underdosing of other psychotropic drugs, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.8X6S	Underdosing of other psychotropic drugs, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T43.96XA	Underdosing of unspecified psychotropic drug, init encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.96XD	Underdosing of unspecified psychotropic drug, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T43.96XS	Underdosing of unspecified psychotropic drug, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T44.0X6A	Underdosing of anticholinesterase agents, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T44.0X6D	Underdosing of anticholinesterase agents, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T44.0X6S	Underdosing of anticholinesterase agents, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T44.1X6A	Underdosing of other parasympathomimetics, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T44.1X6D	Underdosing of other parasympathomimetics, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T44.1X6S	Underdosing of other parasympathomimetics, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T44.2X6A	Underdosing of ganglionic blocking drugs, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T44.2X6D	Underdosing of ganglionic blocking drugs, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T44.2X6S	Underdosing of ganglionic blocking drugs, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T44.3X6A	Underdosing of oth parasympatholytics and spasmolytics, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T44.3X6D	Underdosing of oth parasympatholytics and spasmolytics, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T44.3X6S	Underdosing of parasympatholytics and spasmolytics, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T44.4X6A	Underdosing of predominantly alpha-adrenocpt agonists, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T44.4X6D	Underdosing of predominantly alpha-adrenocpt agonists, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T44.4X6S	Underdosing of predom alpha-adrenocpt agonists, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T44.5X6A	Underdosing of predominantly beta-adrenocpt agonists, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T44.5X6D	Underdosing of predominantly beta-adrenocpt agonists, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T44.5X6S	Underdosing of predom beta-adrenocpt agonists, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T44.6X6A	Underdosing of alpha-adrenoreceptor antagonists, init encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T44.6X6D	Underdosing of alpha-adrenoreceptor antagonists, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T44.6X6S	Underdosing of alpha-adrenoreceptor antagonists, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T44.7X6A	Underdosing of beta-adrenoreceptor antagonists, init encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T44.7X6D	Underdosing of beta-adrenoreceptor antagonists, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T44.7X6S	Underdosing of beta-adrenoreceptor antagonists, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T44.8X6A	Underdosing of centr-acting/adren-neurn-block agnt, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T44.8X6D	Underdosing of centr-acting/adren-neurn-block agnt, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T44.8X6S	Underdosing of centr-acting/adren-neurn-block agnt, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T44.906A	Undrdose of unsp drugs aff the autonomic nervous sys, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T44.906D	Undrdose of unsp drugs aff the autonomic nervous sys, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T44.906S	Undrdose of unsp drugs aff the autonm nervous sys, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T44.996A	Underdosing of drug aff the autonomic nervous system, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T44.996D	Underdosing of drug aff the autonomic nervous system, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T44.996S	Underdosing of drug aff the autonomic nervous sys, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.0X6A	Underdosing of antiallergic and antiemetic drugs, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.0X6D	Underdosing of antiallergic and antiemetic drugs, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.0X6S	Underdosing of antiallergic and antiemetic drugs, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.1X6A	Underdosing of antineoplastic and immunosup drugs, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.1X6D	Underdosing of antineoplastic and immunosup drugs, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.1X6S	Underdosing of antineoplastic and immunosup drugs, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T45.2X6A	Underdosing of vitamins, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.2X6D	Underdosing of vitamins, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.2X6S	Underdosing of vitamins, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.3X6A	Underdosing of enzymes, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.3X6D	Underdosing of enzymes, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.3X6S	Underdosing of enzymes, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.4X6A	Underdosing of iron and its compounds, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T45.4X6D	Underdosing of iron and its compounds, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.4X6S	Underdosing of iron and its compounds, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.516A	Underdosing of anticoagulants, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.516D	Underdosing of anticoagulants, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.516S	Underdosing of anticoagulants, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.526A	Underdosing of antithrombotic drugs, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.526D	Underdosing of antithrombotic drugs, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T45.526S	Underdosing of antithrombotic drugs, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.606A	Underdosing of unsp fibrinolysis-affecting drugs, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.606D	Underdosing of unsp fibrinolysis-affecting drugs, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.606S	Underdosing of unsp fibrinolysis-affecting drugs, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.616A	Underdosing of thrombolytic drugs, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.616D	Underdosing of thrombolytic drugs, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.616S	Underdosing of thrombolytic drugs, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T45.626A	Underdosing of hemostatic drugs, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.626D	Underdosing of hemostatic drugs, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.626S	Underdosing of hemostatic drugs, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.696A	Underdosing of oth fibrinolysis-affecting drugs, init encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.696D	Underdosing of oth fibrinolysis-affecting drugs, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.696S	Underdosing of other fibrinolysis-affecting drugs, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.7X6A	Undrdose of anticoag antagonist, vit K and oth coag, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T45.7X6D	Undrdose of anticoag antagonist, vit K and oth coag, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.7X6S	Undrdose of anticoag antagonist, vit K and oth coag, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.8X6A	Underdosing of primarily systemic and hematolog agents, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.8X6D	Underdosing of primarily systemic and hematolog agents, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.8X6S	Undrdose of primarily systemic and hematolog agents, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.96XA	Undrdose of unsp prim systemic and hematolog agent, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T45.96XD	Undrdose of unsp prim systemic and hematolog agent, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T45.96XS	Undrdose of unsp prim systemic and hematolog agent, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.0X6A	Underdosing of cardi-stim glycos/drug similar act, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.0X6D	Underdosing of cardi-stim glycos/drug similar act, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.0X6S	Underdosing of cardi-stim glycos/drug similar act, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.1X6A	Underdosing of calcium-channel blockers, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.1X6D	Underdosing of calcium-channel blockers, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.1X6S	Underdosing of calcium-channel blockers, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T46.2X6A	Underdosing of other antidysrhythmic drugs, init encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.2X6D	Underdosing of other antidysrhythmic drugs, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.2X6S	Underdosing of other antidysrhythmic drugs, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.3X6A	Underdosing of coronary vasodilators, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.3X6D	Underdosing of coronary vasodilators, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.3X6S	Underdosing of coronary vasodilators, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.4X6A	Underdosing of angiotens-convert-enzyme inhibitors, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T46.4X6D	Underdosing of angiotens-convert-enzyme inhibitors, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.4X6S	Underdosing of angiotens-convert-enzyme inhibitors, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.5X6A	Underdosing of other antihypertensive drugs, init encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.5X6D	Underdosing of other antihypertensive drugs, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.5X6S	Underdosing of other antihypertensive drugs, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.6X6A	Underdosing of antihyperlip and antiarterio drugs, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.6X6D	Underdosing of antihyperlip and antiarterio drugs, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T46.6X6S	Underdosing of antihyperlip and antiarterio drugs, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.7X6A	Underdosing of peripheral vasodilators, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.7X6D	Underdosing of peripheral vasodilators, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.7X6S	Underdosing of peripheral vasodilators, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.8X6A	Undrdose of antivaric drugs, including scler agents, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.8X6D	Undrdose of antivaric drugs, including scler agents, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.8X6S	Undrdose of antivaric drugs, including scler agents, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T46.906A	Underdosing of unsp agents aff the cardiovascular sys, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.906D	Underdosing of unsp agents aff the cardiovascular sys, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.906S	Underdosing of unsp agents aff the cardiovasc sys, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.996A	Underdosing of agents aff the cardiovascular sys, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.996D	Underdosing of agents aff the cardiovascular sys, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T46.996S	Underdosing of agents aff the cardiovascular sys, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T47.0X6A	Underdosing of histamine H2-receptor blockers, init encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T47.0X6D	Underdosing of histamine H2-receptor blockers, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T47.0X6S	Underdosing of histamine H2-receptor blockers, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T47.1X6A	Underdosing of antacids and anti-gastric-secretory drugs, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T47.1X6D	Underdosing of antacids and anti-gastric-secretory drugs, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T47.1X6S	Underdosing of antacids and anti-gastric-secretory drugs, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T47.2X6A	Underdosing of stimulant laxatives, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T47.2X6D	Underdosing of stimulant laxatives, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T47.2X6S	Underdosing of stimulant laxatives, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T47.3X6A	Underdosing of saline and osmotic laxatives, init encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T47.3X6D	Underdosing of saline and osmotic laxatives, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T47.3X6S	Underdosing of saline and osmotic laxatives, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T47.4X6A	Underdosing of other laxatives, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T47.4X6D	Underdosing of other laxatives, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T47.4X6S	Underdosing of other laxatives, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T47.5X6A	Underdosing of digestants, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T47.5X6D	Underdosing of digestants, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T47.5X6S	Underdosing of digestants, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T47.6X6A	Underdosing of antidiarrheal drugs, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T47.6X6D	Underdosing of antidiarrheal drugs, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T47.6X6S	Underdosing of antidiarrheal drugs, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T47.7X6A	Underdosing of emetics, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T47.7X6D	Underdosing of emetics, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T47.7X6S	Underdosing of emetics, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T47.8X6A	Underdosing of agents primarily affecting GI sys, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T47.8X6D	Underdosing of agents primarily affecting GI sys, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T47.8X6S	Underdosing of agents primarily affecting GI sys, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T47.96XA	Underdosing of unsp agents aff the GI sys, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T47.96XD	Underdosing of unsp agents aff the GI sys, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T47.96XS	Underdosing of unsp agents aff the GI sys, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T48.0X6A	Underdosing of oxytocic drugs, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T48.0X6D	Underdosing of oxytocic drugs, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T48.0X6S	Underdosing of oxytocic drugs, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T48.1X6A	Underdosing of skeletal muscle relaxants, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T48.1X6D	Underdosing of skeletal muscle relaxants, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T48.1X6S	Underdosing of skeletal muscle relaxants, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T48.206A	Underdosing of unsp drugs acting on muscles, init encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T48.206D	Underdosing of unsp drugs acting on muscles, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T48.206S	Underdosing of unspecified drugs acting on muscles, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T48.296A	Underdosing of other drugs acting on muscles, init encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T48.296D	Underdosing of other drugs acting on muscles, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T48.296S	Underdosing of other drugs acting on muscles, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T48.3X6A	Underdosing of antitussives, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T48.3X6D	Underdosing of antitussives, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T48.3X6S	Underdosing of antitussives, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T48.4X6A	Underdosing of expectorants, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T48.4X6D	Underdosing of expectorants, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T48.4X6S	Underdosing of expectorants, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T48.5X6A	Underdosing of other anti-common-cold drugs, init encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T48.5X6D	Underdosing of other anti-common-cold drugs, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T48.5X6S	Underdosing of other anti-common-cold drugs, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T48.6X6A	Underdosing of antiasthmatics, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T48.6X6D	Underdosing of antiasthmatics, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T48.6X6S	Underdosing of antiasthmatics, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T48.906A	Undrdose of unsp agents prim acting on the resp sys, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T48.906D	Undrdose of unsp agents prim acting on the resp sys, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T48.906S	Undrdose of unsp agents prim acting on the resp sys, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T48.996A	Underdosing of agents primarily acting on the resp sys, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T48.996D	Underdosing of agents primarily acting on the resp sys, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T48.996S	Undrdose of agents primarily acting on the resp sys, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T49.0X6A	Underdosing of local antifung/infect/inflamm drugs, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T49.0X6D	Underdosing of local antifung/infect/inflamm drugs, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T49.0X6S	Underdosing of local antifung/infect/inflamm drugs, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T49.1X6A	Underdosing of antipruritics, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T49.1X6D	Underdosing of antipruritics, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T49.1X6S	Underdosing of antipruritics, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T49.2X6A	Underdosing of local astringents and local detergents, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T49.2X6D	Underdosing of local astringents and local detergents, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T49.2X6S	Underdosing of local astringents/detergents, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T49.3X6A	Underdosing of emollients, demulcents and protectants, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T49.3X6D	Underdosing of emollients, demulcents and protectants, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T49.3X6S	Underdosing of emollients, demulcents and protect, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T49.4X6A	Underdosing of keratolyt/keratplst/hair trmt drug, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T49.4X6D	Underdosing of keratolyt/keratplst/hair trmt drug, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T49.4X6S	Underdosing of keratolyt/keratplst/hair trmt drug, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T49.5X6A	Underdosing of ophthalmological drugs and preparations, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T49.5X6D	Underdosing of ophthalmological drugs and preparations, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T49.5X6S	Underdosing of oph drugs and preparations, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T49.6X6A	Underdosing of otorhino drugs and preparations, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T49.6X6D	Underdosing of otorhino drugs and preparations, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T49.6X6S	Underdosing of otorhino drugs and preparations, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T49.7X6A	Underdosing of dental drugs, topically applied, init encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T49.7X6D	Underdosing of dental drugs, topically applied, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T49.7X6S	Underdosing of dental drugs, topically applied, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T49.8X6A	Underdosing of other topical agents, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T49.8X6D	Underdosing of other topical agents, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T49.8X6S	Underdosing of other topical agents, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T49.96XA	Underdosing of unspecified topical agent, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T49.96XD	Underdosing of unspecified topical agent, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T49.96XS	Underdosing of unspecified topical agent, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.0X6A	Underdosing of mineralocorticoids and their antag, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.0X6D	Underdosing of mineralocorticoids and their antag, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T50.0X6S	Underdosing of mineralocorticoids and their antag, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.1X6A	Underdosing of loop diuretics, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.1X6D	Underdosing of loop diuretics, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.1X6S	Underdosing of loop [high-ceiling] diuretics, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.2X6A	Underdosing of crbnc-anhydr inhibtr, benzo/oth diuretc, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.2X6D	Underdosing of crbnc-anhydr inhibtr, benzo/oth diuretc, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.2X6S	Undrdose of crbnc-anhydr inhibtr, benzo/oth diuretc, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T50.3X6A	Underdosing of electrolytic/caloric/wtr-bal agnt, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.3X6D	Underdosing of electrolytic/caloric/wtr-bal agnt, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.3X6S	Underdosing of electrolytic/caloric/wtr-bal agnt, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.4X6A	Underdosing of drugs affecting uric acid metabolism, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.4X6D	Underdosing of drugs affecting uric acid metabolism, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.4X6S	Underdosing of drugs affecting uric acid metabolism, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.5X6A	Underdosing of appetite depressants, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T50.5X6D	Underdosing of appetite depressants, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.5X6S	Underdosing of appetite depressants, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.6X6A	Underdosing of antidotes and chelating agents, init encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.6X6D	Underdosing of antidotes and chelating agents, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.6X6S	Underdosing of antidotes and chelating agents, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.7X6A	Underdosing of analeptics and opioid receptor antag, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.7X6D	Underdosing of analeptics and opioid receptor antag, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T50.7X6S	Underdosing of analeptics and opioid receptor antag, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.8X6A	Underdosing of diagnostic agents, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.8X6D	Underdosing of diagnostic agents, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.8X6S	Underdosing of diagnostic agents, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.906A	Underdosing of unsp drug/meds/biol subst, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.906D	Underdosing of unsp drug/meds/biol subst, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.906S	Underdosing of unsp drug/meds/biol subst, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T50.996A	Underdosing of drug/meds/biol subst, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.996D	Underdosing of drug/meds/biol subst, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.996S	Underdosing of drug/meds/biol subst, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.A16A	Undrdose of pertussis vaccine, inc combin w pertuss, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.A16D	Undrdose of pertussis vaccine, inc combin w pertuss, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.A16S	Undrdose of pertussis vaccine, inc combin w pertuss, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.A26A	Underdosing of mixed bacterial vaccines w/o a pertuss, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T50.A26D	Underdosing of mixed bacterial vaccines w/o a pertuss, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.A26S	Undrdose of mixed bacterial vaccines w/o a pertuss, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.A96A	Underdosing of other bacterial vaccines, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.A96D	Underdosing of other bacterial vaccines, subs encntr	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.A96S	Underdosing of other bacterial vaccines, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.B16A	Underdosing of smallpox vaccines, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.B16D	Underdosing of smallpox vaccines, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T50.B16S	Underdosing of smallpox vaccines, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.B96A	Underdosing of other viral vaccines, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.B96D	Underdosing of other viral vaccines, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.B96S	Underdosing of other viral vaccines, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.Z16A	Underdosing of immunoglobulin, initial encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.Z16D	Underdosing of immunoglobulin, subsequent encounter	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.Z16S	Underdosing of immunoglobulin, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

T50.Z96A	Underdosing of oth vaccines and biological substances, init	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.Z96D	Underdosing of oth vaccines and biological substances, subs	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
T50.Z96S	Underdosing of vaccines and biological substances, sequela	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code is for "underdosing" of drugs, when a patient takes less than prescribed. This concept doesn't exist at all in ICD-9. This code represents a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y70.0	Diagnostic and monitoring anesth devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y70.1	Therapeutic and rehab anesth devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y70.2	Prosth/oth implnt/mtrls anesthesiology devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y70.3	Surgical instrumnt, matrl and anesth devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y70.8	Miscellaneous anesthesiology devices assoc w incdt, NEC	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

Y71.0	Diagnostic and monitoring cardiovasc devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y71.1	Therapeutic and rehab cardiovasc devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y71.2	Prosth/oth implnt/mtrls cardiovascular devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y71.3	Surg instrumnt, matrl and cardiovasc devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y71.8	Miscellaneous cardiovascular devices assoc w incdt, NEC	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y72.0	Diagnostic and monitoring otorhino devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y72.1	Therapeutic and rehab otorhino devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y72.2	Prosth/oth implnt/mtrls otorhino devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y72.3	Surgical instrumnt, matrl and otorhino devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

Y72.8	Miscellaneous otorhino devices assoc w incdt, NEC	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y73.0	Dx and monitor gastroent and urology devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y73.1	Theraputc and rehab gastroent and urology dev assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y73.2	Prosth/oth implnt/mtrls gastroent and urol dev assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y73.3	Surg instrumnt, matrl & gastroent and urol dev assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y73.8	Misc gastroent and urology devices assoc w incdt, NEC	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y74.0	Dx and monitoring gen hosp/persnl-use devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y74.1	Theraputc and rehab gen hosp/persnl-use dev assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y74.2	Prosth/oth implnt, matrl & hosp/persnl-use dev assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

Y74.3	Surg instrumnt,matrl & gen hosp/persnl-use dev assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y74.8	Miscellaneous gen hosp/persnl-use devices assoc w incdt, NEC	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y75.0	Diagnostic and monitoring neurological devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y75.1	Therapeutic and rehab neurological devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y75.2	Prosth/oth implants, matrl and neuro devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y75.3	Surgical instrumnt, matrl and neuro devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y75.8	Miscellaneous neurological devices assoc w incdt, NEC	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y76.0	Diagnostic and monitoring ob/gyn devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y76.1	Therapeutic and rehab ob/gyn devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

Y76.2	Prosth/oth implnt/mtrls ob/gyn devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y76.3	Surgical instrumnt, matrl and ob/gyn devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y76.8	Miscellaneous ob/gyn devices assoc w incdt, NEC	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y77.0	Diagnostic and monitoring ophthalmic devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y77.1	Therapeutic and rehab ophthalmic devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y77.2	Prosth/oth implnt/mtrls ophthalmic devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y77.3	Surgical instrumnt, materials and opth devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y77.8	Miscellaneous ophthalmic devices assoc w incdt, NEC	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y78.0	Diagnostic and monitoring radiological devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

Y78.1	Therapeutic and rehab radiological devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y78.2	Prosth/oth implnt/mtrls radiological devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y78.3	Surgical instrumnt, matrl and radiolog devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y78.8	Miscellaneous radiological devices assoc w incdt, NEC	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y79.0	Diagnostic and monitoring orthopedic devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y79.1	Therapeutic and rehab orthopedic devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y79.2	Prosth/oth implnt/mtrls orthopedic devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y79.3	Surgical instrumnt, materials and orth devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y79.8	Miscellaneous orthopedic devices assoc w incdt, NEC	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

Y80.0	Dx and monitoring physical medicine devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y80.1	Theraputc and rehab physical medicine devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y80.2	Prosth/oth implnt/mtrls physical med devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y80.3	Surg instrumnt, matrl and physcl med devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y80.8	Miscellaneous physical medicine devices assoc w incdt, NEC	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y81.0	Dx and monitoring gen/plast-surg devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y81.1	Therapeutic and rehab gen/plast-surg devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y81.2	Prosth/oth implnt/mtrls gen/plast-surg devices assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y81.3	Surg instrumnt, matrl and gen/plast-surg dev assoc w incdt	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

Y81.8	Miscellaneous gen/plast-surg devices assoc w incdt, NEC	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y82.8	Other medical devices associated with adverse incidents	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y82.9	Unsp medical devices associated with adverse incidents	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y90.0	Blood alcohol level of less than 20 mg/100 ml	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y90.1	Blood alcohol level of 20-39 mg/100 ml	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y90.2	Blood alcohol level of 40-59 mg/100 ml	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y90.3	Blood alcohol level of 60-79 mg/100 ml	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y90.4	Blood alcohol level of 80-99 mg/100 ml	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y90.5	Blood alcohol level of 100-119 mg/100 ml	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

Y90.6	Blood alcohol level of 120-199 mg/100 ml	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y90.7	Blood alcohol level of 200-239 mg/100 ml	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y90.8	Blood alcohol level of 240 mg/100 ml or more	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y90.9	Presence of alcohol in blood, level not specified	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Y95	Nosocomial condition	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.00	Weeks of gestation of pregnancy not specified	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.01	Less than 8 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.08	8 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.09	9 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

Z3A.10	10 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.11	11 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.12	12 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.13	13 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.14	14 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.15	15 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.16	16 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.17	17 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.18	18 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

Z3A.19	19 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.20	20 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.21	21 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.22	22 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.23	23 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.24	24 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.25	25 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.26	26 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.27	27 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

Z3A.28	28 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.29	29 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.30	30 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.31	31 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.32	32 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.33	33 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.34	34 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.35	35 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.36	36 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

Z3A.37	37 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.38	38 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.39	39 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.40	40 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.41	41 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.42	42 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z3A.49	Greater than 42 weeks gestation of pregnancy	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z67.10	Type A blood, Rh positive	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z67.11	Type A blood, Rh negative	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).

Z67.20	Type B blood, Rh positive	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z67.21	Type B blood, Rh negative	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z67.30	Type AB blood, Rh positive	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z67.31	Type AB blood, Rh negative	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z67.40	Type O blood, Rh positive	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z67.41	Type O blood, Rh negative	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z67.90	Unspecified blood type, Rh positive	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).
Z67.91	Unspecified blood type, Rh negative	NoDx	This code represents a condition for which there are no corresponding clinical concepts in ICD-9. This code represent a condition that is not likely to require backwards mapping (which is used primarily for financial or actuarial purposes).